

AUGUST 1, 1949

MODERN MEDICINE

The Journal of Diagnosis and Treatment

What Can Be Done
for the Hay Fever
Patient? (see page 47)



Complete table
of contents page 10

Dr. Albert V. Stoesser (see page 10)



A good night's rest
A good day's work

**Allergic patients get both, with
just 4 small doses**

Comfort 'round-the-clock for your allergy patients . . . Decapryn provides long-lasting relief with low milligram dosage. "Symptoms were relieved from 4 to 24 hours after the administration of a single dose of Decapryn—"1

"It was found that 12.5 mg. could be given during the day with comparatively few side reactions and yet maintain good clinical results—"2

prescribe

Decapryn[®] SUCCINATE

Brand of Doxylamine Succinate

THE LONG-LASTING LOW-DOSAGE ANTIHISTAMINE

12.5 mg. tablets, P. R. N. Also available in pleasant tasting syrup especially designed for children. (6.25 mg. per 5 cc) and 25 mg. tablets.



CINCINNATI • U.S.A.

1. Sheldon, J. M. et al: Univ. Mich. Hosp. Bull. 14:13-15 (1948). 2. MacGiddy, E. L.: Neb. State M. J. 34:123 (1949)

DAYTIME SEDATION

BUTISOL[®] SODIUM

BRAND OF EUTABARBITAL SODIUM

"Intermediate Sedative"



Mild in action, without sharp peaks of effect, Butisol Sodium is well suited to daytime sedation. Its action has been described¹ as "intermediate" between the fast-acting derivative, pentobarbital, and the longer-acting barbital and phenobarbital.

Butisol is destroyed in the body—is not dependent upon renal excretion. With proper regulation of dosage there is no cumulative action and a minimum of "hang-over."

DOSAGE FORMS: Elixir Butisol Sodium, 0.2 Gm. (3 gr.) per fl. oz.*—Also Tablets and Capsules, 0.1 Gm. (1½ gr.); Tablets, 15 mg. (¼ gr.), 30 mg. (½ gr.)*, and 50 mg. (¾ gr.).

Caution: Use only as directed.

*Samples on Request



Elixir Butisol Sodium—bright, green color; inviting flavor; excellent prescription vehicle.

McNEIL

LABORATORIES, INC.

PHILADELPHIA 32, PENNSYLVANIA

¹ Council on Pharmacy and Chemistry: J.A.M.A., 135:224. (Sept. 27) 1947.





Psoriasis Circinata Before RIASOL



Psoriasis Circinata After RIASOL

RIASOL



**for
greater
summer
freedom**

in PSORIASIS

Summer heat and humidity frequently aggravate the discomforts of the psoriatic sufferer. The unsightly lesions not only hamper summer freedom, but often cause added embarrassment as well.

Your prescription for RIASOL is doubly appreciated in hot weather, because it affords striking cosmetic relief and a sense of greater freedom in most instances. Recurrences are usually minimal.

RIASOL is safe and effective and may be applied to any part of the body, including the scalp and face. Patients like RIASOL because it is simple, convenient and pleasant to use. No bandages necessary.

RIASOL contains 0.45% mercury chemically combined with soaps, 0.5% phenol and 0.75% cresol in a washable, non-staining, odorless vehicle.

Apply daily after a mild soap bath and thorough drying. A thin, invisible, economical film suffices. After one week, adjust to patient's progress.

RIASOL is ethically promoted. Supplied in 4 and 8 fld. oz. bottles, at pharmacies or direct.

MAIL COUPON TODAY — TEST RIASOL YOURSELF

SHIELD LABORATORIES
12850 Mansfield Ave., Detroit 27, Mich.

MM-8-49

Please send me professional literature and generous clinical package of RIASOL.

..... M.D. Street

City Zone State

Druggist Address



RIASOL FOR PSORIASIS



when reducers stray from the dietary path...

... DESOXYN Hydrochloride provides a safe, simple and effective curb on the wayward appetite. At the same time, the stimulating action of DESOXYN increases the patient's sense of well-being and desire for activity. To depress the appetite, one 2.5-mg. tablet an hour before breakfast and lunch is usually sufficient. A third tablet may be taken in midafternoon, if needed, and if it does not cause insomnia. • It has been shown that weight for weight DESOXYN is more potent than other sympathomimetic amines so that *smaller doses* may be used effectively. In addition, DESOXYN has a *faster action, longer effect and relatively few side-effects*. • Orally, DESOXYN is an effective cerebral stimulant with a wide variety of uses. Parenterally, it helps to restore and maintain blood pressure during surgery. For detailed information, simply drop a card to ABBOTT LABORATORIES, NORTH CHICAGO, ILLINOIS

TABLETS,

2.5 mg. and 5 mg.

ELIXIR,

20 mg. per fluidounce
(2.5 mg. per fluidrachm)

AMPOULES,

20 mg. per cc.

Prescribe

desoxyn[®] HYDROCHLORIDE

(Methamphetamine Hydrochloride, Abbott)

MODERN MEDICINE



THE JOURNAL OF DIAGNOSIS AND TREATMENT

EDITORIAL BOARD

James T. Case, M.D., *Chicago*
Franklin D. Dickson, M.D., *Kansas City*
Julius Hess, M.D., *Chicago*
Walter B. Hoover, M.D., *Boston*
Foster Kennedy, M.D., *New York City*
John C. Krantz, Jr., PH.D., *Baltimore*
A. J. Lanza, M.D., *New York City*
Milton S. Lewis, M.D., *Nashville*
George R. Livermore, M.D., *Memphis*
Francis W. Lynch, M.D., *St. Paul*
Cyril M. MacBryde, M.D., *St. Louis*
Karl A. Meyer, M.D., *Chicago*
Alton Ochsner, M.D., *New Orleans*
Robert F. Patterson, M.D., *Knoxville*
Edwin B. Plimpton, M.D., *Los Angeles*
Fred W. Rankin, M.D., *Lexington, Ky.*
John Alton Reed, M.D., *Washington*
Rufus S. Reeves, M.D., *Philadelphia*
Dalton K. Rose, M.D., *St. Louis*
Howard A. Rusk, M.D., *New York City*
Roger S. Siddall, M.D., *Detroit*
James S. Simmons, M.D., *Boston*
W. Calhoun Stirling, M.D., *Washington*
Frank P. Strickler, M.D., *Louisville*
Richard Torpin, M.D., *Augusta, Ga.*
Paul M. Wood, M.D., *New York City*

SCIENCE WRITERS: Thomas B. Gibbons, M.D.,
William F. Maloney, M.D., Bernardine
Lufkin, Patricia O. Read, Elizabeth
Kane, Edward S. Niederkorn

TRANSLATORS: John W. Brennan, M.D.,
Julius Lichtenstein

EDITORIAL CONSULTANTS

Carolyn G. Adams, M.D.
E. R. Anderson, M.D.
Joe W. Baird, M.D.
Edward P. Burch, M.D.
James B. Carey, M.D.
C. D. Creevy, M.D.
C. J. Ehrenberg, M.D.
W. K. Haven, M.D.
Charles Hymes, M.D.
Miland E. Knapp, M.D.
Ralph T. Knight, M.D.
John F. Pohl, M.D.
Leo Rigler, M.D.
M. B. Sinykin, M.D.
A. V. Stoesser, M.D.
Arthur L. H. Street, LL.B.
Marvin Sukov, M.D.
Harry A. Wilmer, M.D.

EDITORIAL STAFF

A. E. Hedback, M.D.
Editor
Thomas Ziskin, M.D.
Associate Editor
Mark S. Parker
Executive Editor
Sarah A. Davidson
Managing Editor
Daisy Stilwell
Art Editor
James Niess
*Editorial Board
Secretary*
Margaret W. Brodt
Marjorie M. Voigt
Editorial Assistants




DR. VERRIE WYSE SAYS:

*They're so easy to use . . .
so complete and accurate.*

HISTACOUNT

PATIENTS' RECORD FORMS



Note how continuation sheet fits into folded "Histacount" form and how that form can hold other papers.

More than 50,000 doctors use and approve "Histacount" forms. Every year thousands of other doctors "discover" these records. Their overwhelming popularity and general acceptance are their best testimonial.

THE POPULAR FOLDER STYLE

These forms provide twice as much record space as standard cards . . . yet take no more room. When folded (to card size) they hold other data pertaining to your patients.


Choose from more than 245 forms

LETTER-SIZE AND CARD STYLE FORMS

Two more complete lines of "Histacount" forms designed to suit your needs: Letter-size forms to fit regular letter files and the old reliable card forms (5 x 8)

FREE SAMPLES AND CATALOGUE

Actual samples of "Histacount" forms and copy of 48-page catalogue are yours on request.



Letter-Size
8 1/2" x 11"

Card-Size
5" x 8"

**PROFESSIONAL
PRINTING COMPANY, INC.**

America's Largest Printers to the Professions

202 TILLARY ST., BROOKLYN 1, N. Y.

PROFESSIONAL PRINTING CO., INC.
202-208 Tillary St., Brooklyn 1, N. Y.

Please send your 48-page Histacount Catalogue and actual samples of forms.

Dr. _____ Degree _____

STATIONERY • HISTACOUNT PRODUCTS
PRINTING • RECORDS • FILES & SUPPLIES

Good Grip Restored


In arthritis involving the hands, hips or any other joints, restoration of function and diminution of pain are best accomplished by complete systemic rehabilitation.

Darthronol—furnishing the anti-arthritic effects of massive dosage of vitamin D and the nutritional benefits of 8 other vitamins—plays an important role in the rehabilitation of the arthritic.

a ROERIG preparation

DARTHRONOL

FOR THE ARTHRITIC





each
capsule
contains



Vitamin D (Irradiated Ergosterol)	50,000 U.S.P. Units
Vitamin A (Fish-Liver Oil)	5,000 U.S.P. Units
Vitamin C (Ascorbic Acid)	75 mg.
Vitamin B ₁ (Thiamine Hydrochloride)	3 mg.
Vitamin B ₂ (Riboflavin)	2 mg.
Vitamin B ₆ (Pyridoxine Hydrochloride)	0.3 mg.
Niacinamide	15 mg.
Calcium Pantothenate	1 mg.
d-alpha Tocopherol Acetate	2.4 mg.

(Equivalent by biological assay to 3.3 mg. International Standard Vitamin E)

J. B. ROERIG AND COMPANY • 536 Lake Shore Drive • Chicago 11, Illinois



Contents for August 1 1949

VOL. 17, NO. 15



LETTER FROM THE EDITOR.....14

CORRESPONDENCE..... 18

QUESTIONS & ANSWERS..... 24

FORENSIC MEDICINE..... 34

WASHINGTON LETTER..... 40

SPECIAL ARTICLE

What Can Be Done for the Hay Fever Patient?
Albert V. Stoesser..... 47

MEDICINE

Heparin Therapy of Coronary Thrombosis
Leo Loewe and H. B. Eiber..... 51

Reclassified Cardiovascular Rejectees
Paul D. White, Robert L. Levy, William J. Kerr, William D. Stroud, and George K. Fenn..... 52

Primary Pleural Effusion
Joseph R. Kraft..... 53

Exchange Transfusion Therapy
M. Bessis..... 54

Early Diagnosis of Femoral Thrombosis..... 55

ANESTHESIOLOGY

Baird's Anesthesia for Infants
Christine F. Webster and Frederick H. Van Bergen..... 58

THE MAN ON THE COVER against a background of pollen-bearing greenery is Dr. Albert V. Stoesser, director of Allergy Clinics at the Minneapolis General and the University of Minnesota Hospitals, Minneapolis. In addition to his practice he has been teaching at the University of Minnesota Medical School for the past twenty years. A pediatrician whose interests have led him to become an allergist, he is active in several associations of pediatricians, allergists, immunologists, and pathologists. Dr. Stoesser is author of the Special Article, "What Can Be Done for the Hay Fever Patient?" on page 47.



drowsiness *minimized . . .*

allergic patients remain alert . . .

Clinical reports describing the use of Thephorin in 2564 patients with hay fever and other allergies indicate an incidence of drowsiness of only 2.92%. In contrast with other antihistamines, Thephorin can therefore be given to motorists and other patients who have to remain alert. Highly effective and well tolerated in most cases,

Thephorin is available in 25-mg tablets and as a palatable syrup which permits convenient adjustment of dosage.

HOFFMANN-LA ROCHE INC • NUTLEY 10 • N. J.

Thephorin[®]
brand of phenindamine

'Roche'



Contents
for
August 1
1949

CONTINUED

SURGERY

- Prevention of Amputation Neuroma
Stephen Teneff..... 59
- Surgical Treatment of Cardiospasm
Rodney Maingot..... 60
- Pericardial Aspiration..... 62

DERMATOLOGY

- Poison Ivy Management
J. B. Howell..... 63
- When Sunshine Is Harmful
J. M. Hitch..... 66
- Pruritus..... 67
- Dermatophytosis..... 67

PEDIATRICS

- Juvenile Cirrhosis of the Liver
Paul D. Keller and William L. Nute, Jr. 68
- Curare and Exercise for Poliomyelitis
W. D. Paul and O. A. Couch, Jr...... 69

OBSTETRICS

- Heart Disease and Pregnancy
John J. Sampson..... 70

GYNECOLOGY

- Relief from Dysmenorrhea..... 71
- Basal Metabolism during Menopause..... 72

MEDICAL FORUM

- Basket for Removing Ureteral Stones..... 72

DIAGNOSTIX..... 74


SHORT REPORTS..... 81

CURRENT BOOKS & PAMPHLETS..... 96

MEDICAL FILMS..... 100

PATIENTS I HAVE MET..... 102

MODERN MEDICINE, The Journal of Medical Progress, is published twice monthly on the first and fifteenth of each month at Minneapolis, Minn. Subscription rate: \$5.00 a year, 25c a copy. Business Manager: M. E. Herz. Address editorial correspondence to 84 South 10th Street, Minneapolis 3, Minn. Telephone: Bridgeport 1291. ADVERTISING REPRESENTATIVES: New York 17: George Doyle, Bernard A. Smiler, Lee Klemmer, 1 East 42nd Street, Suite 801, Corn Exchange Bank Bldg. Telephone: MUrray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Central 6-4619. SAN FRANCISCO 4: Duncan A. Scott & Co., Mills Bldg. Telephone: Garfield 1-7950. LOS ANGELES 5: Duncan A. Scott & Co., 2978 Wilshire Blvd. Telephone: Dunkirk 8-4151. SEATTLE: Duncan A. Scott & Co., 827 Securities Bldg. Telephone: Seneca 6135.




ARGYPULVIS 98% Effective in TRICHOMONIASIS

... according to published report
of Reich, Button and Nechtow, "Treatment
of *Trichomonas Vaginalis* Vaginitis,"
Surgery, Gynecology and Obstetrics,
May, 1947, pp. 891-896.*



For Use by the Physician
7-gram bottles fitting
Halmespray or
equivalent powder-
blower (in cartons of 3)

For Home Use
by the Patient
2-gram capsule
for insertion
by the patient
(in bottles of 12)



ARGYPULVIS

ARGYROL and ARGYPULVIS are registered trade-
marks, the property of

A. C. BARNES COMPANY
NEW BRUNSWICK, N. J.

• The technique included both office insufflation and supplementary home use of the powder in capsules, along with usual precautions against reinfection. A most significant observation was that results from use of the capsules alone were approximately the same.

Thus, this new form of dependable ARGYROL affords the physician a highly efficient weapon in the control of Trichomoniasis, through office and home treatment or by the patient's home use of capsules alone.

Composition ... Physical Properties
ARGYPULVIS contains powdered ARGYROL (20%), Kaolin (40%) and Beta Lactose (40%) ... finely milled, to provide the fluffiness which makes for easy insufflation, and with an attraction for water which promotes fast action.

INTRODUCTORY TO PHYSICIANS: *On request we will send professional samples of ARGYPULVIS (both forms), together with a reprint of the Reich, Button and Nechtow report. (Use coupon.)

A. C. Barnes Company
Dept. MM-89 New Brunswick, N. J.

Name

Address

City.....State.....

LETTER FROM THE EDITOR

Dear Reader:

Every item you read in MODERN MEDICINE is aimed at a target which we call The Reader. The Reader is a composite of all doctors and is a most demanding gentleman.

He wants accurate information quickly. He wants simple, interesting language without windy rhetoric, and he wants to stop reading when the data has been delivered. He puts up with no nonsense and is a leading factor in development of ulcers among our editorial staff.

We do not always please The Reader, but our efforts to do so are never ending. It is this devotion to a cause that gives distinction to the reports in MODERN MEDICINE.

A change of pace is provided in our Correspondence section. There *the readers* take over. There you have your say and write what you will. You praise, criticize, and condemn. Sometimes you tell about an unusual case or a project close to your heart. Sometimes you ask for advice, sometimes you give it.



One doctor in Chicago was hypersensitive to noise. The issue containing his letter was hardly in the mail before *the readers* picked up their pens. All suggestions were relayed to the Chicago physician, who was most grateful. A similar response greeted the plea of a Florida practitioner with Parkinson's syndrome.

When a Chinese medical library asked for books to replace those destroyed by war, you sent in hundreds of tomes.

Not all contributions have been in a serious vein. The matter of the proper Chinese term for constipation was a challenge. Your response was even more terrific than your puns.



a complete
gallbladder
drug regimen
in a
single tablet



NEOCHOLAN

(A BRAND OF DEHYDROCHOLIC ACID COMP., PHENOBARBITAL AND HOMATROPINE METHYLBROMIDE, P.-M. CO.)

FOR NONCALCULOUS BILIARY-TRACT STASIS

Formula: Each tablet of Neocholan contains:

Dehydrocholic Acid Comp., P.-M. Co.....	265 mg. (4 grs.)
Dehydrocholic Acid.....	250 mg. (3 3/4 grs.)
Phenobarbital.....	.80 mg. (1/8 gr.)
Homatropine Methylbromide.....	1.2 mg. (1/50 gr.)

Rationale: Neocholan exerts two distinct actions:

1 . . . Hydrocholeresis—the dehydrocholic acid content stimulates the liver to secrete a greatly increased flow of thin, free-flowing bile, thereby tending to flush the biliary ductal system.

2 . . . Sphincter Relaxation—homatropine methylbromide and phenobarbital combine to assure sphincter relaxation so that the bile may discharge freely into the duodenum.

Neocholan, therefore—used in conjunction with appropriate dietary measures— provides the complete gallbladder regimen in a single tablet.

For increased efficiency, ease of administration and economy—

R NEOCHOLAN

SUPPLIED IN BOTTLES OF 100 TABLETS

PITMAN MOORE COMPANY

PHARMACEUTICAL AND BIOLOGICAL CHEMISTS
DIVISION OF ALLIED LABORATORIES INC., INDIANAPOLIS 6, INDIANA

AN IMPORTANT NEW MILESTONE in the Therapy of Arthritic Affections

The "cure of [rheumatic fever]", agree most authoritative sources,^{2,3,5,6} "depends not only on reaching, but also on maintaining a high plasma salicylate level."⁵ The correlation between such blood levels and symptomatic improvement is graphically shown in the table at the right.⁶

Pabalate—latest product of Robins' research—now helps to achieve and maintain *higher salicylate blood levels on lower salicylate dosage.*

This is made possible by the combination in Pabalate of non-toxic para-aminobenzoic acid with sodium salicylate. As visualized in the chart at the lower right,⁶ para-aminobenzoic acid (itself an active antirheumatic)⁷ manifests a reciprocal action with salicylates when administered concurrently—sharply increasing the blood salicylate levels (under constant salicylate dosage),^{1,4} and in turn having its own blood levels effectively enhanced.⁴

The clinical significance of this synergistic relationship represents an important advantage in the therapy of arthritic affections. Pabalate 'Robins'—a strictly ethical preparation—is available at (or may be secured by) all leading pharmacies.

A. H. ROBINS CO., INC., RICHMOND 20, VA.
Ethical Pharmaceuticals of Merit since 1878

indications: Rheumatoid arthritis; acute rheumatic fever; fibrositis; gout; osteo-arthritis.

dosage: Two or three enteric coated tablets every three or four hours, without sodium bicarbonate.

formula: Each enteric coated tablet contains:

Sodium salicylate, U.S.P. (5 gr.), 0.3 Gm.;

Para-aminobenzoic acid (as sodium salt), (5 gr.), 0.3 Gm.

supplied: In bottles of 100 and 500 enteric coated tablets.

Salicylate blood level in mg. per 100 cc.	Erythrocyte sedimentation rate
less than 20 mg.	no fall
20-30 mg.	slow fall
30-40 mg.	RETURN TO NORMAL CLINICAL SYMPTOMS SUBSIDE



**For high salicylate blood levels
on low salicylate dosage—**

Pabalate®

Pabalate tablets are enteric coated
to prevent gastric irritation and
insure optimal toleration.




ref
1. B
77:
2. Br
and
3. Co
He
4. Dr
Me
5. Ed
6. Re
17:
7. No
Pro
68:



references:

1. Bellale, M.: Union Med. Can., 77:392, 1948.
2. Brodie, B. B. et al.: J. Pharmacol. and Exper. Therap., 80:114, 1944.
3. Coburn, A. F.: Bull. Johns Hopkins Hosp., 78:458, 1948.
4. Dry, T. J. et al.: Proc. Staff Meetings Mayo Clin., 51:497, 1946.
5. Editorial: J.A.M.A., 138:267, 1948.
6. Reid, J.: Quarterly J. Med., 17:139, 1948.
7. Rosenblum, H. and Fraser, L. E.: Proc. Soc. Exper. Biol. and Med., 65:178, 1947.



Correspondence

Communications from the readers of MODERN MEDICINE are always welcome. Address communications to The Editors of MODERN MEDICINE, 84 South 10th St., Minneapolis 3, Minn.

Cyanosis from Contaminated Water

TO THE EDITORS: In the Refresher in General Practice on "Care of the Newborn" (*Modern Medicine*, Mar. 15, 1949, p. 36), I note the omission of the possibility of nitrite-producing organisms in the intestinal tract as a contamination from the water used in making up formulas, causing either met- or sulfhemoglobinemia. This can be the cause of cyanosis, easily confirmed (and cured) by changing to proper supply of water.

PAUL W. VAN METRE, M.D.
Rockwell City, Iowa

More Special Articles

TO THE EDITORS: Please print more special articles and exhibits.

CHOW CHAN, M.D.
Hong Kong, China

Always Reads "Forensic Medicine"

TO THE EDITORS: I read with a great deal of interest your column on Forensic Medicine each time in my copy of *Modern Medicine*. It is contributing to my better understanding of legal medicine. I have formed the habit of clipping out certain interesting legal points which some day may be of interest to me professionally as well as academically.

ROBERT B. ELLIOTT, M.D.
Houston

Cover to Cover

TO THE EDITORS: I like your magazine very much and it is the only one I read regularly from cover to cover.

C. E. WILLS, JR., M.D.
Washington, Ga.

Physicians' Letter Exchange

TO THE EDITORS: The United Nations Council of Philadelphia has been working for some time on the promotion of international friendships by a Letter Exchange between America and the other countries of the world.

There has been a tremendous response from people in all walks of life and we feel sure that professional men, and physicians in particular, could help tremendously by exchanging ideas with foreign colleagues.

American physicians, returning from abroad, tell of many discouraging conditions—lack of sufficient drugs and equipment and, most important, ignorance of the advances in various fields of medicine during the past ten years.

Foreign physicians are eager for correspondence with American colleagues. A British physician writes that he is anxious to "exchange views with an un-nationalized doctor." We have just received lists of hundreds of Japanese medical men and students



A NEW, DRAMATIC THERAPY FOR THE RELIEF OF PAIN AND LESIONS OF

HERPES ZOSTER

DESCRIPTION: Protamide is a sterile, aqueous colloidal solution of a specially processed proteolytic enzyme, for the maximum relief of nerve root pains of Herpes Zoster and Tabes Dorsalis.


CLINICAL RESULTS: Highly gratifying clinical results have been obtained with the use of Protamide (Sherman) in the treatment of the extremely resistant herpes syndrome. Pain has been relieved in the great majority of herpes cases within four to forty-eight hours and lesions have healed in ten days or less—regardless of the particular nerve roots involved. Complete clinical data may be obtained by writing for the Protamide literature on Herpes Zoster and a recent reprint on Protamide for Tabes Dorsalis.

DOSAGE: In Herpes Zoster the recommended dosage is 1.3 cc of Protamide intramuscularly each day from two to four days. No contraindications or incompatibility have been reported to date. All Protamide is clinically tested for positive results. Can be stored at room temperature without loss of potency.

**ALSO CLINICALLY PROVED
FOR THE LIGHTNING PAINS
AND ATAXIA
OF TABES DORSALIS**

SHERMAN LABORATORIES
G. H. Sherman, M. D., Founder
BIOLOGICALS • PHARMACEUTICALS
DETROIT 15, MICHIGAN

*
REGISTERED U. S. TRADE MARK



THE SHOE
that lives up to your
recommendations...

JUMPING-JACKS®
FLEXIBLE SHOES FOR HARD WEAR

Pediatricians and doctors throughout the U. S. and many foreign countries enthusiastically recommend Jumping-Jacks for children six months to four years of age. Jumping-Jacks' unique, patented, one-piece sole and heel helps prevent ankles from turning—gets little feet off to a good start in life.



VAISEY-BRISTOL SHOE CO., INC.
ROCHESTER 3, NEW YORK
MONETT, MISSOURI • SKOWHEGAN, MAINE
MADE IN CANADA BY THE
SAVAGE SHOE COMPANY, LTD. • PRESTON, ONTARIO

interested in discussing all branches of medicine. What better method of establishing international friendships than by exchanging ideas and discussing conditions with a foreign friend?

A physician can help by writing to one foreign physician. Each correspondent will be matched according to age, background, and, as near as possible, field of medicine. The foreign letters will be translated whenever necessary.

Letters should be addressed to: Letters Abroad, United Nations Council of Philadelphia, 1411 Walnut Street, Philadelphia 3.

G. ALISON RAYMOND
Publicity Director

Philadelphia

An Efficient Physician's Bag

TO THE EDITORS: Upon starting practice following the war, it soon became apparent to me that the usual physician's bag is a very inefficient gadget. Perhaps many doctors have shared my experience of groping unsuccessfully in the dark recesses of the bag for an item they cannot find and finally in desperation turning the bag upside down. Usually, the last item to fall out is what you're looking for and that breaks as it hits the floor.

Following such an especially trying search I started working on a different style of bag and after several unsuccessful tries, happened to see a mechanic's tool kit which was modified by a cabinetmaker to reduce its size. This was, however, too big. I wrote to the company who made this bag and tried to interest them in making a physician's bag. They made up a bag to my specifications and after using it for some time it was further modified. The photo shows the final product.



Familiar to every physician are the



patients who need a good tonic



To stimulate appetite, to restore vigor and general tone, Eskay's NEURO PHOSPHATES and Eskay's THERANATES are two of the most useful preparations you have. These tonics are prescribed so widely because they work so well.

Smith, Kline & French Laboratories,
Philadelphia

Eskay's Neuro Phosphates

a palatable and effective tonic

Eskay's Theranates

the formula of famous Neuro Phosphates, plus Vitamin B₁

Flavettes

in hypertension

A NEW
BENIGN
COMPOUND
TO CURB THE APPETITE

FLAVETTES have been found to be effective in curbing the appetite and securing weight loss in 80% of 568 cases¹ regardless of the clinical indication and specifically in hypertension where thyroid and amphetamine are contraindicated. FLAVETTES is not an "obesity tablet", but a product capable of appeasing the appetite rather than oxidizing excessive weight. Samples and literature on request.

1. Gould, W. L.: New York
St. J. Med. 47:981, 1947

AMHERST RESEARCH, INC.
Box 3503 Merchandise Mart Station
Chicago 54, Ill.

The bag consists of an upper compartment which runs the whole length and width of the bag and is 3 in. deep. Pill bottles are attached to the cover. The lower portion of the bag consists of five square drawers of varying depth. The drawers have small grooves in their sides for the insertion of metal partitions in either a lateral or longitudinal direction. The ends of the drawers are covered when the bag is closed and the cover locks into place when the bag is closed. The bag is made of wood and the bottom of the drawers are metal. The outside is covered with conventional black leather and the top edges are rounded.

By being square the space in the



bag is used more efficiently than if the top were rounded as the conventional bag is. The drawers are compartmentalized to accommodate syringes, needles, ampules, and instruments, while larger objects can be carried in the top compartment. The bag weighs 8 1/2 lb., empty. Outside dimensions are: 11 1/2 in. high, 8 in. wide, and 14 in. long.

A. T. HAEREM, M.D.
Redwood City, Calif.

5

good reasons

There are five good reasons why SULAMYD* (Sulfacetimide-Schering) is the preferred sulfonamide in the treatment of pyelitis, pyelonephritis, cystitis and other infections of the urinary tract.



SULAMYD

(SULFACETIMIDE-SCHERING)

1

Unusual efficacy: In *B. coli* infections of the urinary tract, recovery or improvement with SULAMYD is unusually high, ranging from 93 to 98% of cases.^{1,2}

2

High urinary levels: Due to SULAMYD's rapid clearance from the blood and easy excretion by the kidneys, high urinary concentrations are obtained in the presence of low blood levels.³

3

Excellent renal tolerance: Crystalluria is rare and anuria has never been reported following SULAMYD therapy. This enviable record is due, in large measure, to SULAMYD's remarkable solubility (946 mg. per 100 cc.), approximately 80 times that of sulfadiazine.³

4

Systemically well tolerated: Side effects are minimal and uncommon. SULAMYD may, therefore, be administered with greater security to infants, children and pregnant women.^{1,4}

5

Alkalinisation is unnecessary, since SULAMYD is quite soluble in both acid and alkaline urine.

PACKAGING: SULAMYD, sulfacetimide, tablets of 0.5 Gm., in bottles of 100 and 1,000 tablets; and bottles of 5.0 Gm. powder, for laboratory determinations upon blood and urine.

BIBLIOGRAPHY: (1) Welebir, F., and Barnes, R.W.; J.A.M.A. 117:2132, 1941. (2) Alyea, E.P., and Parrish, A.A.; South. M.J. 40:678, 1947. (3) Lehr, D.; J. Urol. 54:87, 1945. (4) Prentiss, R.J., and Kanealy, J.F.; J. Urol. 87:31, 1962.

*S

Schering

CORPORATION-BLOOMFIELD, NEW JERSEY
IN CANADA, SCHERING CORPORATION LIMITED, MONTREAL

SULAMYD



Questions & Answers

All questions received will be answered by letter directed to the petitioner; questions chosen for publication will appear with the physician's name deleted. Address all inquiries to the Editorial Department, MODERN MEDICINE, 84 South Tenth Street, Minneapolis 3, Minnesota.

QUESTION: I have a twenty-nine-year-old man suffering—and I do mean suffering—from migratory phlebitis. Numerous TT's have been used to no avail. Available literature gives little encouragement. Could you offer some therapeutic suggestions?

M.D., Pennsylvania

ANSWER: By *Consultant in Vascular Diseases*. Many believe that migratory phlebitis is an early symptom of Buerger's disease. I treat my cases with hot packs locally and large doses of penicillin. I use the fortified penicillin, 300,000 units every forty-eight hours. To this I often add 7.5 gr. of sulfadiazine four times a day. When the attack is completely over I continue the sulfadiazine for one week and then for one week in every four during the next year. The patient should stop smoking.

INSTRUCTIONS FOR HOT PACKS

Bed rest is essential. It is useless to sit in a chair with the leg on another chair and try to apply hot wet packs.

Body must be flat. Only one pillow under head.

Knee should be flexed slightly with the lower leg high and about level on a pile of pillows 16 in. above the mattress.

Packs should be applied at least 10 in. above and below the area to be treated.

Temperature of packs should begin at 120° F. and be increased as tolerated—preferably to 130° F. if the patient does not object.

Fresh packs should be applied as

quickly as possible to prevent cooling of the skin between changes. The packs are best applied as follows:

- 1] Fold a blanket four times and place on top of a pile of pillows.
- 2] Half fill several hot-water bottles with water at right temperature. Compress the air out of the bottles as cork is tightened.
- 3] Wring large bath towels and woolen blankets out of hot water. Quickly apply the wet towels, then the hot-water bottles, then wrap the blanket about all, and pin tightly.

If the pack is properly applied it will stay hot for at least two hours. The pack should be changed every two hours.

Best results will be obtained if the packs are continued for not less than four hours at a time.

When the ACE bandage is applied the patient should walk a great deal.

QUESTION: How much cyclopropane is used for an anesthesia of one-half hour, as for routine appendectomy? How much ethylene?

M.D., Tennessee

ANSWER: By *Consultant in Anesthesiology*. The question cannot be answered in exactly this form. If the purpose of the question is to find the total amount of gas used and therefore the cost, the information may be figured approximately from the following:

Usually the 5-liter bag is filled with 80% oxygen and 20% cyclopropane. The mask is fitted tightly and the gases are set at about 300 cc. per minute each

TYMPANIDE



Contains:

ETHYLENE GLYCOL Monostearate
fungicide and bactericide

SULFANILAMIDE 5% bactericide

UREA 10% cleansing bactericide

ISOBUTANOL 3% topical anesthetic

WARREN-TEED

Pharmaceuticals

THE WARREN-TEED COMPANY • COLUMBUS 8 • OHIO

Meyenberg

evaporated
GOAT MILK

Gives prompt
proven relief



FOR THE colic, diarrhea or vomiting of cow's milk lactalbumin allergy...or in borderline cases when such sensitivity is suspected, prescribe Meyenberg, the original evaporated goat milk.

Meyenberg Evaporated Goat Milk is nutritionally equivalent to evaporated cow's milk—economical, sterilized, easy to prepare. Available at all pharmacies in 14-oz. hermetically-sealed containers.



SPECIAL MILK PRODUCTS, INC.

LOS ANGELES 25



CALIFORNIA

of oxygen and cyclopropane. The cyclopropane is increased by 100 cc. per minute every thirty seconds until flowing at 600 cc. per minute. The cyclopropane is kept at that rate until the proper plane of anesthesia is attained, usually seven or eight minutes after the start of anesthesia. From then on the cyclopropane flow usually ranges between 50 cc. and 200 cc. per minute for correct anesthesia.

For ethylene the 5-liter bag may be filled with 80% ethylene and 20% oxygen. The flow may then be continued with 200 cc. ethylene and 300 cc. oxygen with a tight-fitting mask and enough ether vapor added to obtain the right anesthesia.

QUESTION: Will you give me some information regarding the technic for performing the Rotter test for tissue levels of vitamin C? Where can I obtain the dichlorophenol-indophenol solution? How do I interpret the results?

M.D., Oklahoma

ANSWER: By Consultant in Clinical Pathology. Since the description of the intradermal dye discoloration test for vitamin C by Rotter in 1938, several articles have appeared in which various authors have attempted evaluations (L. B. Slobody, *J. Lab. & Clin. Med.* 29:464, 1944; A. H. Holland et al., *J. Lab. & Clin. Med.* 32:124, 1947).

Slobody's procedure is to inject enough $\frac{N}{300}$ 2-6 dichlorophenol-indophenol intradermally to raise a wheal 4 mm. in diameter. The time of complete disappearance of the dye is measured. Slobody interprets the results as follows:


Complete decolorization in over 14 minutes—Vitamin C deficiency

Complete decolorization in 9 to 13 minutes—Mild deficiency

Complete decolorization in less than 9 minutes—No deficiency



*Psoriasis — duration
15 years*



*Photographed after 8 weeks
treatment with Mazon*

Mazon and Mazon Soap act quickly and bring skin infections under control*

**These infections include*

Eczema
Psoriasis
Alopecia
Dandruff
Ringworm
Ivy Poisoning
Athlete's foot

The fundamental principles employed in the development of Mazon—careful selection of ingredients, skillful blending for ease of application, absorbability and patient acceptance—promote the successful treatment of obstinate skin conditions.

The distinctive characteristics and therapeutic value of Mazon have been demonstrated clinically for over 25 years. Skin irritations not caused by or associated with systematic or metabolic disease or disorders are usually amenable to treatment with Mazon.

MAZON

BELMONT LABORATORIES,
Philadelphia, Pa.

60,000
50,000 DOCTORS
prefer the
HYFREATOR

... for the removal of skin growths, tonsil tags, cysts, small tumors, superfluous hair, and for other technics by electrodesiccation, fulguration, bi-active coagulation.

Now, completely redesigned the new **HYFREATOR** provides more power and smoother control . . . affording better cosmetic results and greater patient satisfaction. Doctors who have used this new unit say it provides for numerous new technics and is easier, quicker to use.

Send for descriptive brochure, "Symposium on Electrodesiccation and Bi-Active Coagulation" which explains the **HYFREATOR** and how it works.



THE BIRTCHER CORPORATION

To: The **BIRTCHER** Corp., Dept. A-8-9
5087 Huntington Dr., Los Angeles 32, Calif.
Please send me free booklet, "Symposium on
Electrodesiccation & Bi-Active Coagulation."

Name _____
Street _____
City _____ State _____

The dye can be obtained from Eastman Kodak Co., Rochester, N. Y., under the name "Sodium 2-6 dichlorobenzenoneindophenol."

The vitamin C saturation test performed by measuring the amount of vitamin C excreted in the urine following an intravenous test dose of vitamin C is considered by most to be a more satisfactory test for vitamin C deficiency.

Complete details of this test can be found in an article by I. S. Wright *et al.*, *Arch. Int. Med.* 60:264, 1937; and one by S. L. Smith, *J.A.M.A.* 111: 1753, 1938.

QUESTION: What difference in cochlear pathology is there between injury from persistent acoustic trauma and that from a single instance of acoustic trauma?

M.D., New York

ANSWER: By Consultant in Otology. In persistent acoustic trauma the pathologic changes are most pronounced in the basal turn of the cochlea. Degeneration occurs in the hair cells of Corti's organ and the cells of the spiral ganglion. There is early loss of hearing in the region of 4,096 cycles.

Later this loss will extend below this level to involve the speech frequencies. This finding is characteristic of progressive inner ear deafness.

Following acoustic trauma of the blast or concussion type, two types of injury are found. In one the middle ear is damaged, with traumatic rupture of the tympanic membrane and hemorrhage. The cochlea may not be seriously harmed. In the second form, the force is exerted mainly on the inner ear, without drum injury. In this case, the pathologic changes are located chiefly in the basal turn of the cochlea just within the oval window.



call it

dermatophytosis pedis
epidermophytosis
tinea pedis
dermatomycosis
trichomycosis
or "athlete's foot"

then
treat it
with

'Timofax'

BRAND

undecylenate
ointment

"Timofax" Ointment makes possible more efficient management of "athlete's foot". It has proved to be remarkably effective—while affording the blandness essential for prolonged prophylaxis and treatment.

Chemically related to the normal constituents of sweat, undecylenic acid does not break down the skin in the manner of a foreign substance. Hopkins et al. state that "The high percentage of effectiveness and low incidence of irritation indicated that undecylenic acid was the best of the fungicides tested for routine treatment of sub-acute cases."

(*J. Invest. Dermat.*, 7:239, 1946)

Timofax


undecylenate
ointment

"Timofax" contains undecylenic acid 10% as free acid and potassium undecylenate in a scented vanishing cream base. Supplied in ¼ oz. collapsible tubes.



BURROUGHS WELLCOME & CO. (U.S.A.)
INC. Tuckahoe 7, N.Y.

BLOOD REGENERATION



IN SECONDARY ANEMIA

- Non-utilization of iron prescribed for the anemic patient, and its consequent excretion fecally, produces gastrointestinal upset and often defeats the original purpose for which it was intended.

Through the catalytic action of the copper sulphate in Copperin, the amount of iron ammonium citrate per capsule is reduced to only 32 mgm., but ALL the iron is made available for regenerative processes and the maximum therapeutic effect is obtained.

Available for adults in Copperin "A"; for children in Copperin "B".

MYRON L. WALKER COMPANY, Inc.
Mount Vernon, New York

*Liberal professional
samples sent
on request*

WATER
SOLUBLE

COPPERIN

NON-
STAINING



"A&D"

nipple care

prevents fissures

"Our observations clearly indicate that the use of [White's] vitamin A and D ointment in the local care of the puerperal nipple gave protective and therapeutic results much better than those obtained by other methods."*

White's Vitamin A and D Ointment keeps the integument soft and free from excessive or abnormal dryness, eliminates the use of alcohol or other astringent medication, hastens healthy granulation and epithelization in fissure therapy. Provides the natural vitamins A and D in a pleasantly fragrant lanolin-petrolatum base.

*Brougher, J. C.: Prevention and Treatment of Postpartum Fissured Nipples with Local Applications of Vitamin A and D Ointment, Western J. Surg., Obstet. and Gynecol. 52:520-521 (Dec.) 1944.

VITAMIN
White's **A&D** **for nipple care**
OINTMENT

1.5 oz. tubes; 8 oz. and 16 oz. jars; 5 lb. containers.

WHITE LABORATORIES, INC., Pharmaceutical Manufacturers, NEWARK 7, N. J.



SET to SPECIALIZE




Progressive doctors who specialize in eye, ear, nose and throat work, as well as general practitioners, recognize the advantages in owning a modern diagnostic set. They know how it helps alleviate their burden and speeds up examination and operative time. Bausch & Lomb offers both the Arc-Vue Prism Otoscope and the improved May Ophthalmoscope as a unit attractively cased. You owe it to yourself as well as your patients to have modern equipment for fast, accurate diagnoses.

BAUSCH & LOMB

OPTICAL COMPANY



ROCHESTER 2, N. Y.



irradiated !

to
prevent
homologous
Serum
Hepatitis !

Ultraviolet irradiation of plasma destroys not only bacteria but also viral contaminants that might cause homologous serum hepatitis. • You may therefore administer irradiated LYOVAC plasma without danger of serum hepatitis from the plasma infusion. • Stable, portable LYOVAC Normal Human Plasma (*Irradiated*) is prepared from fresh, citrated, human blood of healthy donors, according to regulations of the National Institute of Health. The plasma is pooled, flash frozen, *dehydrated from the frozen state under high vacuum* (lyophile process), then sealed under vac-

uum. • *A blood substitute of choice for emergencies*, irradiated LYOVAC plasma is quickly restored, needs no typing or cross-matching, and each unit is osmotically equivalent to two units of whole blood. • LYOVAC Normal Human Plasma (*Irradiated*) is supplied in vacuum bottles to yield 50 cc., 250 cc. and 500 cc. of restored, irradiated normal human plasma, (containing 660 mgm. of *gamma globulin* in each 100 cc.) or smaller quantities of concentrated, *hypertonic* plasma, for special purposes. Sharp & Dohme, Philadelphia 1, Pa.

Lyovac®

SHARP
& DOHME

Normal Human Plasma IRRADIATED

Forensic Medicine

COMPILED BY ARTHUR L. H. STREET, LL.B.

PROBLEM: Was an employee entitled to workmen's compensation award for disability resulting from smallpox vaccination supplied by the employer, when the vaccination was urged, but not required?

COURT'S ANSWER: Yes.

The New Jersey Superior Court, Appellate Division, said that the disabling effects of the vaccination should be regarded as an accident arising out of and in the course of employment; that it would be unrealistic to assume that the employer furnished the vaccination solely for the benefit of his employees. The vaccination benefited the employer by acting as a safeguard against the serious effects of smallpox among his employees (64 Atl. 2d 99).

PROBLEM: Staff members of a memorial hospital contracted to render services for fixed percentages of money received by the hospital from members and nonmembers. Did it constitute an illegal waste of the hospital's funds for the directors to pay a bonus of \$1,500 in a lump sum to each of the doctors comprising the staff?

COURT'S ANSWER: No.

The U. S. Court of Appeals, Tenth Circuit, upholding a decision of the U.S. District Court for Kansas, noted that the bonus was paid in recognition of efficient services rendered over a four-year period and in compliance with an agreement that if the percent-

age compensation should prove to be inadequate during the early years of the hospital, a bonus would be paid if financial success was achieved later. The court said that a bonus paid by a corporation in good faith for services is not ordinarily to be regarded as a reckless expenditure or unwarranted gift if it is related to the value of services rendered and promotes the corporation's best interests (170 Fed. 2d 859).

PROBLEM: A newspaper published a letter stating that a doctor had refused to attend an accident victim because he was not licensed to practice in the state. The paper commented editorially that if an out-of-state doctor could not legally render first aid in an emergency, the law was silly and dangerous and that in such emergencies a doctor ought to be permitted to render first aid without technically violating the law. Did the newspaper defame the doctor?

COURT'S ANSWER: No.

The New Hampshire Supreme Court decided that the letter and comment were not so worded as to show intention of the publisher to reflect discredit upon the doctor. The jury decided that the editor merely intended to condemn the law if it forbade emergency treatment by a doctor not locally licensed. The Supreme Court ruled that the verdict was supported by ample evidence. The Supreme Court noted that, although local law

**individuality
in
spasmolytic
prescription
writing**

Individualized management of complex spastic syndromes is particularly facilitated by use of Donnatal Elixir—the spoonfed spasmolytic agent whose superior efficacy derives from its precise balance of the principal *natural* alkaloids of belladonna, plus phenobarbital. • With it may be administered judicious selections from a wide variety of gastric, intestinal, bronchial, urinary, analgesic, or other drug agents with which it is entirely compatible for the precise pharmacotherapy you desire.

Write for suggestive formulary for your ready reference.

Each Sec of Donnatal Elixir contains:

Hyoscyamine Sulfate	0.1037 mg.
Atropine Sulfate	0.0194 mg.
Hyoscine Hydrobromide	0.0065 mg.
Phenobarbital (1/4 gr.)	16.2 mg.

Also available as Donnatal Tablets and Donnatal Capsules

**for
spasmolysis
plus**

donnatal elixir
®

Also available as
Donnatal Tablets
and Donnatal Capsules



A. H. Robins Co., Inc.

Ethical Pharmaceuticals of Merit since 1878

Richmond 20, Va.



pain!...

*the synergic
formula
for maximum
non-narcotic
analgesia*

To obtund pain without recourse to narcotics—yet better than the patient's medicine cabinet can—becomes a daily professional obligation. That's why Phenaphen was formulated with calculated pharmacologic precision... the analgesic action of its aspirin—phenacetin components being implemented and prolonged by its phenobarbital content (which helps allay apprehension... its hyoscyamine further increasing overall efficiency through local anodyne action. Phenaphen—the astute professional prescription for pain—is promoted to physicians only.

Each tablet or capsule contains:

Phenacetin (3 gr.)	194 mg.
Acetylsalicylic Acid (2½ gr.)	162 mg.
Hyoscyamine Sulfate	0.03 mg.
Phenobarbital (¼ gr.)	16.2 mg.

● *phenaphen*



A. H. Robins Co., Inc.

ETHICAL PHARMACEUTICALS OF MERIT SINCE 1878

Richmond 20, Va.

did permit the doctor to give emergency aid, he had given the newspaper reason to believe that it did not. The court also upheld the right of newspapers to discuss and condemn in good faith and without malice, as a matter of social and moral duty, the inability of a community to provide emergency care (63 Atl. 2d 791).

PROBLEM: An injured workman brought a malpractice suit against a physician chosen, under the Maine workmen's compensation act, to treat him. Could the workman sue without first demanding that his employer bring the suit?

COURT'S ANSWER: No.

Under the Maine statutes, an injured employee's acceptance of an award does not preclude him from suing the physician, if the employer fails to do so within thirty days after written demand by the employee. Demand upon the employer to sue, however, is a condition to the employee's right to sue.

In common-law personal injury suits, defendant is liable for aggravation of the injury due to malpractice of a physician selected with reasonable care. The increased injury is regarded as a foreseeable result of the original injury.

The Maine Supreme Judicial Court decided that the rule is also applied in workmen's compensation cases when an injury to an employee is aggravated by the negligent or unskillful treatment of a properly chosen physician or surgeon and if the chain of causation remains unbroken the resulting disability or death is compensable and an award of compensation includes the original injury and the ultimate results through malpractice (63 Atl. 2d 302).

An important and pleasant adjunct to sulfonamide therapy




BiSoDoL's quick-acting, pleasant-tasting formula provides the kind of long-lasting alkalization the sulfonamides demand. The proven effectiveness of this outstanding antacid alkalizer merits your professional consideration. Try BiSoDoL next time you prescribe sulfonamide therapy.

BiSoDoL



POWDER
MINTS

WHITEHALL PHARMACAL COMPANY
22 EAST 40TH STREET, NEW YORK 16, N. Y.



VALUE IN NERVE-DEAFNESS ESTABLISHED



A growing number of clinical reports establishes that AMVITOL* (parenterally) supplemented by HYVANOL* (orally) increases hearing acuity in a high proportion of cases of nerve-deafness. Jacobson,¹ a pioneer in the use of vitamin-amino acid therapy in hearing disorders, reported that improvement was shown for the most part in the higher octaves, which he regards as of especial significance since "it is the high-frequency range which is first affected in damages to the acoustic nerve." Later studies² confirmed these original findings: improvement was noted in 73 per cent of 57 patients. Gordon³ obtained favorable results in 76 per cent of 28 cases, using AMVITOL and HYVANOL. Hearing improvement was obtained "within several weeks" after the beginning of treatment. Other investigators^{4,5} have similarly acclaimed this effective, new approach to the management of nerve-deafness.

Because the AMVITOL-HYVANOL method of therapy is free from distressing side-effects, the cooperation of the patient is easily secured. AMVITOL and HYVANOL are available through all prescription pharmacies. Complete literature to physicians on request.

1. Jacobson, M.: New York State J. Med. 45: 2079 (1945).

2. Hirschfeld, H.; Jacobson, M., and Jelinek, A.: Arch. Otolaryngol. 44: 686 (1946).

3. Gordon, G. R.; J. M. A. Alabama 17: 340 (1948).

4. Michels, M. W.; et al.: Permanente Foundation M. Bull. 5: 124 (1947).

5. Laub, G. R.: The Recorder 11: 10 (1947).

*Exclusive trademark of Walker Vitamin Products, Inc.

Walker VITAMIN PRODUCTS, INC., MOUNT VERNON, N.Y.


Ithyphen in

Obesity

Effective Factor

in Weight Reduction

In the treatment of Obesity and in the reducing problem, the administration of ITHYPHEN is of important note. Surprising satisfactory results are obtained from this product without causing acidosis, artificial hypothyroidism or other undue toxic by-effects.



**STRAUSS
LABORATORIES**

**1328 Broadway
NEW YORK 1 N. Y.**

Write for literature
and physician's
sample. Kindly
enclose professional
card or letterhead.

**SUCCESSFULLY
PRESCRIBED
OVER
19
YEARS**

**CONSISTENT
RESULTS WITH
NO STARVATION
OR APPETITE
REDUCTION**

Washington Letter

Survey of Diagnostic Centers

Preliminary reports on a nationwide survey of public nonprofit diagnostic clinics show some surprising results. U.S. Public Health Service experts have established that there is a great variation in the efficiency and effectiveness of the centers. Experts also learned that on the part of physicians there is a similar variation in reliance on these laboratories.

The result is a tremendous, continuous waste of skilled manpower. Because inefficient and ineffective procedures in some clinics produce a high percentage of confusing reports, many

physicians suspect the findings of all diagnostic laboratories. To correct this, U.S. Public Health Service has started a campaign to standardize laboratory methods and procedures. This will be followed by an educational campaign designed to reach every public and nonprofit institution in the country. The expectation is that both tasks can be completed in the next three years.

National Health Council Approves Rapid Expansion

For several years the Communicable Disease Center at Atlanta has been studying clinic procedures. The director, Dr. Seward Miller, presented the findings at a recent meeting of the National Advisory Health Council in Washington. He did not make a formal written report, but the figures he gave were enough to convince the council that need for corrective action was urgent.

The council advised Public Health Service to expand the program as rapidly as possible. No difficulty is anticipated in getting the necessary money for the campaign. Six represent-

(Continued on page 44)



"You make an excellent cup of coffee, Miss Blanchard."



Contact

OF A SENSITIZED BODY CELL
with an allergen and subsequent release of
histamine is considered to be the mechanism
of allergic disorders.

Relief

WITH BENADRYL

BENADRYL, blocking the action of
histamine, prevents reaction in cells that
have been sensitized. Relief of symptoms is
gratifyingly rapid, usually occurring
within an hour or two after the first dose.
And treatment with BENADRYL is simple,
convenient, and inexpensive.



BENADRYL[®]

BENADRYL has been found highly effective in a wide variety of allergic states, ranging from seasonal, such as hay fever, to the non-seasonal, such as acute and chronic urticaria, angioneurotic edema, vasomotor rhinitis, contact dermatitis, erythema multiforme, pruritic dermatoses, dermatographism, serum sickness, food allergy, and sensitization to drugs, such as penicillin and the sulfonamides.

BENADRYL hydrochloride (diphenhydramine hydrochloride, Parke-Davis) is available in a variety of forms to facilitate individualized dosage and flexibility of administration, including Kapseals[®], Capsules and a palatable Elixir.

The usual dosage of BENADRYL is 25 to 50 mg. repeated as required. Children up to 12 years of age may be given 1 to 2 teaspoonsful of Elixir Benadryl.

PARKE, DAVIS & COMPANY • DETROIT 32, MICHIGAN





RESMICON^{T.M.}

acid-adsorbent demulcent

without side effects

BECAUSE

Resmicon combines anion-exchange polyamine resin with natural gastric mucin. This combination means double-barrelled action for relief of pain and promotion of healing in peptic ulcer.

BECAUSE

Resmicon's polyamine resin inactivates hydrochloric acid by physical adsorption, not by chemical "neutralizing" or "buffering." This physical action induces no alkalosis, no acid rebound, no depletion of body electrolytes, no formation of renal calculi, no toxic effects, no irritation, and no uncomfortable side effects such as constipation, diarrhea, eructation, or flatulence.

*for effective
peptic ulcer therapy*



BECAUSE

Resmicon's mucin provides a dense, tenacious coating for the eroded area, as well as for the whole gastric mucosa. This mucinous film—highly resistant to penetration by HCl and pepsin—protects the ulcerated tissue and fosters healing.

RESMICON tablets: gastric mucin, 170 mg.; anion-exchange polyamine resin, 500 mg.; supplied in bottles of 84 tablets.

Whittier
LABORATORIES
Division Nutrition Research Laboratories
Chicago 30, Illinois



WASHINGTON LETTER

atives are in the field gathering material from clinics, and a staff of 80 is at work in Atlanta evaluating the methods and procedures. Next year, when the estimations have been completed and procedures set up, Congress will be asked for an appropriation to cover two years.

Findings Hard to Evaluate

"The first thing we confirmed was that there is a great deal of variety in the effectiveness and methods employed by clinics," a Public Health spokesman told MODERN MEDICINE. "Many of the clinics are not too statistical in their findings, and their reports are difficult to evaluate. It's understandable that doctors don't place the same credence in the findings of every clinic. Some physicians will let findings from a certain clinic influence their decisions 90%, and other men won't give them more than 10% weight. Some take everything in the clinic reports as the Gospel, and others hardly trust them. We think the answer is somewhere in between. But the first step is to develop some uniform procedures in the clinics themselves."

Four Stages to Nationwide Program

The workers at Atlanta are part way through the first phase of the campaign, the "methodology research." Once enough facts have been gathered about how individual laboratories and clinics are doing their work now, standardized methods and procedures will be worked out for all.

In the next phase, key employees of state health departments will be trained in the standardized methods. It will be the task of these state em-

ployees, assisted by the Public Health Service, to train city and county clinic workers in the new methods. Another part of the program will be the development of a uniform diagnosis reference, to insure that the physician and laboratory use and understand specific terms in the same way. Once all this has been accomplished, Public Health Service hopes to confine itself to an advisory role. It will assist states in consultations and evaluations at individual laboratories.

New Cabinet Department by Default?

Unless something unexpected occurs, a new cabinet department of welfare and education will become effective on August 19, with the present Federal Security Administrator, Oscar Ewing, as first secretary. The unexpected development would be adjournment of Congress prior to that date.

The contest against time developed this way: The reorganization act under which President Truman created the new department provides that his recommendations become law if not vetoed by Congress within sixty days. To give himself all the time possible, Mr. Truman sent his reorganization plan to Congress just as soon as he signed the reorganization act, June 20. Early in the session, Congress had been heading for an August 1 adjournment, but that date had to be given up. Now the administration is expecting to keep the session going until after August 19.

The chance that Congress will turn down Mr. Truman by a formal vote is slight. Either house could do this alone and kill the plan, but only by

(Continued on page 78)



Building stones

Tissue repair is the keystone of the recovery process. It makes little difference if the infection is halted, the fracture reduced, or the metabolic imbalance adjusted—it is the patient's own cells that must complete the cure.

While true hypoproteinemia is comparatively rare, nevertheless hypernutrition with essential amino acids during the recovery

process has been shown empirically to speed the patient upon the road to normal health. Amino acid preparations should be supplemented by moderate amounts of vitamins.

Lederle research has for some time been concerned with such mixtures of amino acids and vitamins and their application in the field of nutrition.

LEDERLE LABORATORIES DIVISION

AMERICAN *Cyanamid* COMPANY

30 ROCKEFELLER PLAZA • NEW YORK 20, N. Y.

WYETH INCORPORATED, Philadelphia 3, Pa.



A Complete Line of
MALE AND FEMALE HORMONES

WYETH

Outstanding for Potency, Efficacy, Toleration

TESTOSTERONE	The most potent androgen
CONESTRON® <i>Conjugated Estrogens, Equine, Wyeth</i>	Orally potent natural estrogen for action without side action
PROGESTERONE	Corpus luteum hormone
WYNESTRON*	Aqueous suspension of pure crystalline estrone, Wyeth

All injectable forms now available in the unique, exclusive TUBEX.®

Forms of issue:

Testosterone. Injection Testosterone Wyeth—aqueous suspension of microcrystals of free testosterone for intramuscular injection: multiple dose vials of 10 cc., 10 and 25 mg. per cc.; and vials of 5 cc., 50 mg. per cc. *Membrettes*, Transmucosal Tablets, 4 mg. each: bottles of 30, 100, 500 and 1000. *Implantation Pellets*, 75 mg. each: vials of 1. *Tubex* of 10 and 25 mg., packages of 1 and 5.

Conestron. Tablets of 0.625 and 1.25 mg. (expressed as sodium estrone sulfate): bottles of 100 and 1000.

Progesterone. Injection, aqueous suspension: multiple dose vials of 10 cc., 10 mg. and 25 mg. per cc. *Tubex* of 10 mg., packages of 1 and 5. *Membrettes* of 10 mg., bottles of 100.

Wynestron. Multiple dose vials of 10 cc., 2 mg. per cc. *Tubex* of 1, 2, and 5 mg., packages of 5.

* Trade Mark



MODERN MEDICINE

Special Article

What Can Be Done for the Hay Fever Patient?

ALBERT V. STOESSER, M.D., Ph.D.* †

Prepared for Modern Medicine

MISCONCEPTIONS about hay fever are almost as tenacious as its misnomer, which has persisted for well over a hundred years.

Long ago physicians recognized that the disease was not confined to the haying season and that fever was not necessarily a symptom. Over the past thirty years they have gradually realized that adults are not the only victims, that children, even small infants, may be susceptible.

Much research has been done in the field, many excellent new preparations are offered for therapy. Yet confusion still prevails. Why do so many patients feel that their physicians cannot help them? Why is there so much pessimism about hay fever treatment among doctors and hay fever sufferers?

It may be stated that in spite of all the new preparations offered to the physician, the treatment of pollenosis with drugs alone is not the best. Specific pollen therapy is safer, but even this does not give perfect results. A combination of the two forms of treatment can be recommended. However, greater knowledge of the chemistry of pollen and attention to even minor details in inoculations and related factors are needed.

* Clinical Professor of Pediatrics and Allergy, University of Minnesota Medical School, Minneapolis.

† From the Allergy Clinics, Minneapolis General and University Hospitals, University of Minnesota Medical School, Minneapolis.

SPECIAL ARTICLE

In order to clarify the situation and, we hope, inject a bit of optimism, the problem must be considered from several angles. First, that of diagnosis. Why is the disease so easily overlooked, particularly in children?

Second, what should be done for the hay fever patient in the acute stage of the disease—at the time when he is usually first seen by the physician?

Finally, and most important of all, what are the steps to be taken in the long-range treatment of the disease? Here lies one of the major misunderstandings about hay fever: Too many patients are considered as having a disease which causes trouble for only a few weeks or months of the year. *If every hay fever sufferer were considered a perennial patient, more persons would be benefited by treatment.*

DIAGNOSIS OF HAY FEVER

The onset of hay fever in small children does not always follow a pattern as typical as that in older individuals.

When the causative pollen season starts, which may be at any time during the spring, summer, or fall, the child may seem to have an upper respiratory infection, with or without a little fever. Coughing may appear and, by examination, the clinician becomes suspicious of bronchitis or pneumonia. Recently it has been a too common practice to diagnose the condition, especially if fever is present, as atypical or virus pneumonia.

The results are the same regardless of the kind of therapy tried by different physicians. After a few weeks, the child re-

TABLE 1. SIDE EFFECTS OF ANTIHISTAMINES

<i>Drowsiness</i>	<i>Tremor</i>	<i>Dryness of mouth</i>	<i>Diarrhea</i>
<i>Headache</i>	<i>Muscular weakness</i>	<i>Tachycardia</i>	<i>Constipation</i>
<i>Fatigue</i>	<i>Syncope</i>	<i>Bradycardia</i>	<i>Frequency</i>
<i>Ataxia</i>	<i>Blurred vision</i>	<i>Arrhythmia</i>	<i>Dysuria</i>
<i>Vertigo</i>	<i>Tinnitus</i>	<i>Epigastric pain</i>	<i>Dermatitis</i>
<i>Nervousness</i>	<i>Cough</i>	<i>Nausea</i>	<i>Pruritus</i>
<i>Insomnia</i>	<i>Dyspnea</i>	<i>Vomiting</i>	<i>Anemia</i>

covers. All is well until the following year, when the same sequence occurs. This time the clinician should realize that he is dealing with an allergic manifestation. Most significant is a history of allergy in the family. If the mother or father has eczema, hay fever, or asthma, the diagnosis is practically assured.

The preschool child may have an irritation of the eyes thought to be "pink eye." The first time the trouble is observed, the true cause may not be suspected, but when the condition appears the next year at practically the same time, allergy must be investigated. Tree pollen is the most common offender, but grass or weed pollen may cause the reaction. Many of these patients have no obvious symptoms besides the eye condition, but examination often reveals definite evidence of allergy in the nasal passages.

TABLE 2. RELATIONSHIP OF VARIOUS ANTIHISTAMINIC DRUGS

<i>Group 1</i>		<i>Group 2</i>	
Neohetramine (Wyeth)		Thenylene (Abbott)	
Syrup; 25- 50- and 100-mg. tablets		25- 50- and 100-mg. tablets	
Neo-Antergan (Merck)		Histadyl (Lilly)	
25- and 50-mg. tablets		Syrup; 25- 50- and 100-mg. pulvules; also injectable	
Pyribenzamine (Ciba)		Diatrin (Warner)	
Elixir; 25- and 50-mg. plain tablets; 50-mg. coated tablets		50-mg. tablets	
		Tagathen (Lederle)	
		25-mg. tablets	
		Chlorothen (Whittier)	
		25-mg. tablets	
<i>Group 3</i>		<i>Group 4</i>	
Trimeton (Schering)		Decapryn (Merrell)	
Elixir; 25-mg. tablets		Syrup; 12.5- and 25-mg. tablets	
Pyrrolazote (Upjohn)		Benadryl (Parke, Davis)	
Elixir; 25- and 50-mg. tablets		Elixir; 25- and 50-mg. capsules; also injectable	
Thephorin (Roche)		Hydryllin (Searle)	
Syrup; 25-mg. tablets		Elixir; tablets	

The child of school age may have severe nasal distress during the summer months. Quite often the difficulty begins in the spring and continues into the fall.

Parents often assume that the youngster is having "summer colds," due in part to too much activity in sports during the vacation period; swimming is frequently blamed for the symptoms. However, when the fall of the year arrives and the child

SPECIAL ARTICLE

is back in school, the condition still exists. Not until the following year do further observations bring forth the correct diagnosis.

In the young adult, onset of hay fever is usually more or less typical. The eyes suddenly become injected with lacrimation. Itching is present. The nose is red from rubbing. A rather profuse, watery nasal discharge appears. Sneezing is common. The patient is a truly sick individual, although he has no fever. What shall be done?

IMMEDIATE TREATMENT

The hardest part of the immediate treatment is to convince the adult patient and the parents of the child with hay fever that certain rules are of great importance. How many will cooperate? The number depends chiefly on the amount of time the physician is willing to give to details.

A complete history and physical examination should be obtained in order to get information on any concomitant diseases and to ascertain the patient's general condition.

The diet should be light, but all essential foods must be eaten. Occasionally, frequent small meals are best. Rest is most important. Too much activity for a child or too much work for an adult may greatly increase the severity of symptoms. Long, tiresome rides in the country are forbidden. Swimming is completely eliminated. Dusty, smoky, or even damp places should be avoided.

These recommendations are often disregarded in spite of the fact that they definitely reduce suffering and complications. If adult patients would choose the hay fever season as their vacation period, they might be able to follow the rules more thoroughly.

Medication does not include the routine use of eye or nose drops, although in a few instances this form of therapy may be employed, especially at bedtime. An isotonic solution of 0.25 to 0.5% neosynephrin, 0.5 to 1% propadrine hydrochloride, 1% paredrine hydrobromide, 1 to 2% tuamine sulfate, or 0.05% privine may be tried. Oenethyl sulfate in a 1% solution is a new preparation.

Antistin in the form of a 0.5% solution may relieve eye symptoms but can cause an irritation. The same is true of 0.5% his-

(Continued on page 88)

Heparin Therapy of Coronary Thrombosis

LEO LOEWE, M.D., AND H. B. EIBER, M.D.*

Jewish Hospital, Brooklyn

THE management of patients with acute coronary artery thrombosis should include the use of anticoagulants. Because of easy administration, prompt effectiveness, and lack of toxicity, heparin in Pitkin menstuum is well suited for treatment of the disease. Leo Loewe, M.D., and H. B. Eiber, M.D., feel that heparin in Pitkin menstuum is preferable to dicumarol for avoiding thromboembolic complications in patients with coronary thrombosis.

Because the individual response to dicumarol is variable, the dosage schedule is complex. Dicumarol requires at least twenty-four hours for therapeutic effectiveness and, therefore, offers no protection to the patient during the critical first day of illness. When heparin is given in Pitkin menstuum, prolongation of clotting begins in one to two hours and lasts forty-eight hours or more.

Neither renal nor hepatic disease, though both are contraindications to dicumarol, proscribes the employment of heparin.

Only the relatively simple bedside determination of clotting time is necessary with heparin. When dicumarol is used prothrombin time must be calculated daily, a rather expensive laboratory procedure often difficult to obtain.

Earlier preparations of heparin in

Pitkin menstuum caused a painful local reaction at the site of injection. This effect is now controlled by careful buffering of the gel.

Treatment should be begun as soon as possible after diagnosis is established. For prompt and adequate prolongation of coagulation, 400 mg. of heparin is given subcutaneously in the initial dose of menstuum. This amount is usually repeated every other day for three or four injections. Thereafter, depending on the clotting time, which should be three times normal value or more, the dose of heparin may be diminished to 300 or 200 mg. and given at longer intervals.

Therapy must be continued at least until the patient is allowed out of bed and should then be gradually cut to avoid the hypercoagulability phenomenon which occasionally occurs with sudden cessation of treatment.

If the coagulation time requires shortening, absorption of heparin can be slowed by applying ice bags to the site of the deposit. Circulating heparin is readily inactivated by small transfusions of relatively fresh whole blood.

Active bleeding is the only contraindication to the use of heparin. With an intact vascular system, overdosage causes no complications. The menstuum without vasoconstrictor drugs should be employed when dealing

* Anticoagulation therapy with heparin/Pitkin menstuum in the management of coronary artery thrombosis and its complications. *Am. Heart J.* 37:701-719, 1949.

with intraarterial clotting, especially of the coronary arteries.

Of 20 patients with acute coronary thrombosis treated by heparin in Pitkin menstroom, all but 1 person recovered. Treatment in the fatal case was not begun until the eleventh day of

illness. In the 19 patients who recovered no clinical or electrocardiographic evidence was found of extension of infarct. Periods of bed rest and convalescence were shortened. No thromboembolic complications occurred after anticoagulant therapy.

Reclassified Cardiovascular Rejectees

PAUL D. WHITE, M.D., BOSTON, ROBERT L. LEVY, M.D., NEW YORK CITY, WILLIAM J. KERR, M.D., SAN FRANCISCO, WILLIAM D. STROUD, M.D., PHILADELPHIA, AND GEORGE K. FENN, M.D., CHICAGO*

A STUDY was made in 1947 of 303 men who were reclassified as fit for military service in 1943 after being rejected for circulatory or heart disorders. By assessing the effects of war and postwar adjustment on these men, Paul D. White, M.D., and colleagues sought answers to many problems in cardiovascular diagnosis.

Nearly all the veterans were found to be in good health in 1947. Only 4 were disabled. Most of the men had served long and well. The majority had been on active duty for about two and one-half years and were discharged without disability.

The men were examined by almost exactly the same special cardiovascular boards in Boston, Chicago, New York, Philadelphia, and San Francisco that had reclassified them as 1A in 1943. The total number of men resubmitted was 863, of whom 447 were accepted for military duty. At the time of the study, 303 could be located for examination, all but 49 of whom were veterans.

Heart murmurs, commonest cause for original rejection and present in half of the men in 1947, were usually unimportant and variable.

Transient hypertension, the most difficult condition to evaluate, was observed in 67 men in 1943 and in 17 of these in 1947. Normal readings were noted in 33 of the 67 and sustained high blood pressure in the other 17.

Neurocirculatory asthenia developed in 7 men during the four-year period. Definite heart enlargement was found in only 4 of the veterans. Transient tachycardia, noted in 35 in 1943, had disappeared in 19 by 1947. In a few cases however, tachycardia had developed.

Many of the men had had rheumatic fever in childhood or early adult life, but showed no sequelae. Valvular lesions not evident in 1943 were noted in 4.

* Cardiovascular rejectees. J.A.M.A. 139:1049-1053, 1949.

Primary Pleural Effusion

JOSEPH R. KRAFT, M.D.*

Fitzsimons General Hospital, Denver

UNTIL proved otherwise, primary pleural effusions are usually considered to be tuberculous.

By observation of 100 patients with primary pleural effusion but who did not have parenchymal disease, Joseph R. Kraft, M.D., found pulmonary tuberculosis in 21 and renal tuberculosis in 1 within six months after onset of pleurisy. Other studies have shown that appearance of parenchymal lesions may be expected up to five years after the effusion occurs.

Problems of differentiation are presented by the insidious onset, chest pain, fever, and malaise. The diagnosis of atypical pneumonia with effusion is the most common error.

Unfortunately, no generally applicable diagnostic methods are available for ready identification of the different forms of primary pleurisy.

Examination of aspirated fluid may give definitive information and establish the diagnosis. Nevertheless, the relatively low incidence of successful isolation of the tubercle bacilli from pleural fluid emphasizes the importance of careful physical examination, observation, and judgment from experience.

The treatment of pleural effusions comprises bed rest and removal of sufficient quantity of fluid to limit



the formation of fibrinous visceral-parietal adhesions.

For the probably tuberculous pulmonary lesion the shortest period of hospitalization to be considered is one year of complete bed rest in a sanatorium.

Bed rest for the underlying tuberculous lesion may be considered prophylaxis rather than treatment.

Periodic physical and roentgenographic examinations should be made during four years after discharge from institutional care.

A normal erythrocyte sedimentation rate within the first six months of rest does not necessarily preclude the subsequent development of a parenchymal lesion, and a normal sedimentation rate is not a trustworthy guide by which to determine the duration of treatment.

In many cases, however, subsidence of the erythrocyte sedimentation rate appears to be directly related to regression of pleural fluid or fibrinous pleurisy.

The responsibility upon the physician to regard a pleural effusion as probably tuberculous and to institute appropriate measures is tremendous. A long period of bed rest is a great economic burden. The physician must deliberately disregard this influential factor in making the diagnosis and prescribing the necessary treatment.

* The diagnostic problem of primary pleural effusion. *Am. Rev. Tuberc.* 59:259-269, 1949.

Exchange Transfusion Therapy

M. BESSIS, M.D.*

National Center of Blood Transfusions, Paris

TOTAL replacement transfusion should be considered for severe toxemia when the injurious agent is in the blood, and for leukemia.

M. Bessis, M.D., believes that the most important indication for replacement transfusion is anuric nephritis.

Simultaneous bloodletting and transfusion combats anemia and replaces pathologic plasma with normal plasma, thus preventing or diminishing secondary renal injuries from the products of hemolysis, whether the cause is septicemia, hemolytic poison,

TABLE 1

<i>Replacement Transfusion</i>	<i>Intraperitoneal Dialysis</i>
1. Removes all toxic products, including the nondialysable ones, such as hemoglobin, myoglobin, stromas.	1. Removes only the dialysable toxic products.
2. Does not modify the equilibrium of the tissue fluids.	2. Unless special precautions are taken, equilibrium is destroyed because too many electrolytes are added or lost or too much water added. This may cause cerebral edema.
3. Does not cause any severe reactions.	3. Usually causes peritonitis of the plastic type by the formation of adhesions. Sometimes causes the infectious type.
4. Can be used as often as needed.	4. Possibility of peritonitis prevents continuous use for more than a few days and frequently prevents reemployment.
5. Very efficient. Removes a larger quantity of toxic products which can be calculated beforehand.	5. Removes a smaller quantity which cannot be calculated beforehand.
6. Painless and rapid.	6. Inconvenient and slow.

The procedure provides elimination and allows survival until the kidneys regain normal function.

The effects of replacement transfusion are compared to those of intraperitoneal dialysis in Table 1.

* The use of replacement transfusion in diseases Blood 4:324-337, 1949.

crush syndrome, or transfusion reaction due to incorrect grouping or typing.

Exchange transfusion for leukemia is based on the theory that normal persons have an antileukemic substance other than hemolytic disease of the newborn.

TABLE 2

Quantity of blood injected	Percentage of blood transfused in the patient's organism
1/2 vol. of patient's blood.....	39.4
1 vol.....	63.2
1 1/2 vol.....	77.7
2 vol.....	86.5
2 1/2 vol.....	91.8
3 vol.....	95.0

stance in their blood. Of 38 patients treated at the National Center of Blood Transfusions, Paris, 30 had remissions, 15 with normal peripheral blood, and 6 with normal blood and bone marrow.

Withdrawal and injection of blood are done simultaneously so that total volume is unchanged. The percentage of transfused blood in the body as compared to the quantity injected is shown in Table 2.

For an adult with a blood volume of about 5 liters, 15 liters of the same ABO and Rh groups are needed. If enough of the proper A or B group is not available, O blood may be used after the anti-A and anti-B agglutinins have been neutralized with Wittebsky's AB substances.

Usually 30 donors are necessary. The blood is collected in bottles containing citrate solution.

Serious reactions are prevented if 2 mg. of heparin per kilogram of body weight is given.

Blood is withdrawn from the patient through a plastic catheter in a superficial vein either after cutdown or through a large-bore needle, with the tip pushed up into a larger vein.

An electric pump of Dausset and Moulinier is connected to the catheter and to the donor's blood flask and the used blood receptacle. Blood is drawn at any desired speed, usually about 300 cc. in five minutes. The engine is then reversed to transfuse the patient with donor's blood through the same route. The maneuver is repeated until the desired number of liters is given. Usually an hour is required for children and from two to four hours for adults.

Chills and urticaria occur more frequently with exchange transfusions than with ordinary transfusions, but are not serious. Occasionally the patient's temperature is increased for a day or so.

Replacement transfusions of moderate amounts, about 5 liters, repeated every second or third day, withdraw sufficient urea to enable the patient to survive the critical stage of anuric nephritis. Three months is usually necessary for complete recuperation.

EARLY DIAGNOSIS OF FEMORAL THROMBOSIS is facilitated by the presence of three dilated veins over the tibia. Gerald H. Pratt, M.D., of New York University, New York City, has observed the "sentinel veins" in more than 80% of patients with pathologic clotting and believes the dilated vessels constitute the earliest objective sign of deep vein thrombosis. The sentinel veins are small, superficial, and unsupported by musculature and therefore dilate long before the main veins.

J.A.M.A. 140:476-477, 1949.



The introduction of the New Improved Biolac has met with wide professional interest. Made available as a product of the latest and the most modern refinements in manufacturing facilities, this prescription favorite is *now better than ever.*

NUTRITIONALLY RELIABLE: More than ever a complete food (when vitamin C is added), the New Improved Biolac meets every nutritional requirement of the infant.

All essential fatty acids — with the volatile fraction held to a minimum — are provided by moderate amounts of especially combined fats.

Vegetable and milk sugars — for more satisfactory absorption — are supplied by Biolac's carbohydrate content. Further carbohydrate supplementation is unnecessary.

In protein content, the New Improved Biolac is significantly higher than that of human milk, yielding small, easily digested curds, less allergenic than those of untreated cow's milk.


Prophylactically high levels of such important mineral factors as iron, calcium and phosphorus are incorporated in the New Improved Biolac, together with vitamins A, B₁, B₂ and D. Infant caloric requirements, too, are fully met by Biolac's 20 calories per fluid ounce in standard dilution.

PHYSICALLY IMPROVED: A higher and more stable degree of emulsification of the New Improved Biolac — thereby facilitating digestion — has been brought about by the utilization of every refinement, and the most modern equipment, known to modern infant food manufacturing.

Preparation for feeding is easily calculated; quickly completed — 1 fl. oz. New Improved Biolac to 1½ fl. oz. water per pound of body weight.

NOW, BETTER THAN EVER! The New Improved Biolac can be used interchangeably with the former Biolac which has the same percentage composition of nutritional factors . . . When you prescribe the New Improved Biolac (it costs no more) you may do so with complete confidence. Available only in drugstores, in cans of 13 fl. oz.

THE BORDEN COMPANY
PRESCRIPTION PRODUCTS DIVISION
350 MADISON AVENUE • NEW YORK 17, N. Y.



"Baby Talk for a
Good Square Meal"

Write for professional literature

**THE NEW
IMPROVED BIOLAC**

Baird's Anesthesia for Infants

CHRISTINE F. WEBSTER, M.D., AND FREDERICK H. VAN BERGEN, M.D.*

University of Minnesota, Minneapolis

IN general, the difficulties encountered with the use of anesthetics in infants are similar to those met in adult patients. However, the smaller size of the infant reduces the margin of error and special precautions are necessary.

The infant's high metabolic rate increases the oxygen demand in relation to the small respiratory system. With normal respiration the 21% oxygen content of room air is adequate. When a gauze mask is used for the open drip method of anesthesia only 16% oxygen is provided. This leads to hypoxia.

A closed or semiclosed system permits the administration of sufficient oxygen. To provide the necessary margin of safety, 35% oxygen should be delivered to the patient. To do this and allow for dilution by gas expired from the lungs, oxygen should comprise 50% of the fresh gases entering the closed system.

Even a small mask increases the dead air space, which in the infant represents a much larger percentage of the tidal volume than in the adult. The infant's breathing becomes deep and labored, and exhaustion may quickly appear. Respiratory dead air space may be halved by tracheal intubation and a patent airway assured.

Christine F. Webster, M.D., and Frederick H. Van Bergen, M.D., find

* Pentothal-curare mixture with endotracheal N₂O and O₂ in infants. Bull. Univ. Minnesota Hosp. 20:525-533, 1949.

that many problems are obviated by using Baird's solution for induction and equal parts of nitrous oxide and oxygen by tracheal tube for maintenance of anesthesia. The procedure requires a physician anesthesiologist.

A reliable intravenous route must be maintained, usually requiring preoperative surgical exposure of a vein.

Baird's solution, a mixture containing 25 mg. of pentothal and 5 units of curare (d-tubocurarine chloride) per cubic centimeter, is given in 0.5-cc. doses through intravenous tubing using a three-way stopcock every three minutes until the patient falls asleep. Then 100% oxygen is given by mask and Baird's solution continued until breathing is almost entirely diaphragmatic.

Intubation is then performed, using a Magill oral tube No. 00, 0, or 1, which is 10.5 to 12 cm. in length. The largest diameter easily inserted is employed. Trauma from the procedure is a possibility, but in 27 infants anesthetized by this method laryngeal edema occurred in only 3 cases and was slight. The tracheal tube is connected to a semiclosed system using soda lime to remove carbon dioxide.

Nitrous oxide and oxygen are given at the rate of 500 cc. per minute. Respiration may be regulated and augmented by manual control of the respirator bag. Little additional Baird's

solution is required and the anesthesia is discontinued well before surgery is completed to assure return of reflexes before the tracheal tube is removed.

At the end of the operation the nitrous oxide is washed from the lungs

by giving 100% oxygen. Secretions are aspirated from mouth, pharynx, and trachea, and the tube is removed.

A solution of $\frac{1}{3}$ of normal saline and $\frac{2}{3}$ of 5% glucose is given intravenously throughout the operation.

Prevention of Amputation Neuroma

STEPHEN TENEFF, M.D., ITALY*

PAIN in an amputation stump is usually of nervous origin. Neuroma develops at the severed nerve end when neurofibrils are compressed and strangled by connective tissue. If healing is delayed or if the wound has been infected, the neuroma may be imbedded in the cicatricial tissue of the stump.

Implantation of the cut nerve end into nearby muscles prevents terminal scarring and formation of neuroma and enables the neurofibrils to make physiologic connections with the muscle fibers.

Nerve implantation was employed by Stephen Teneff, M.D., of the University of Turin, Italy, in 6 cases of low- or mid-thigh amputation. Stumps were painless in every instance. In each case after separation of the muscle fibers for a few millimeters, the sciatic nerve was placed unsutured into the body of the adjacent muscle, usually the biceps femoris.

Experiments in rabbits indicate that ten days after operation the nerve stump becomes securely attached to the muscles at the implantation site. Within about twenty-five days, neurofibrils penetrate the narrow zone of loose scar tissue in all directions, separate into fasciculi, and filter into the muscle fasciculi and between the loose connective and intrafibrillar tissue.

From sixty-five to seventy days after operation, the fasciculi of neurofibrils separate and single neurofibrils become firmly attached to the muscle fibers, without trace of connective tissue proliferation.

* Prevention of amputation neuroma. *J. Internat. Coll. Surgeons* 12:16-20, 1949.



Surgical Treatment of Cardiospasm

RODNEY MAINGOT, M.D.*

Royal Free Hospital, London

An esophagus deformed by persistent cardiospasm may be restored to natural shape by incising the constriction down to the mucosa.

Abdominal approach and a variation of Heller's method offer the easiest, safest, and most satisfactory technic, finds Rodney Maingot, M.D., and brought excellent results in 37 of 39 cases not benefited by dilatation. Alternative operations are esophagogastrostomy or esophagocardioplasty.

Habitual spasm of the gastric cardiac sphincter often begins with emotional strain and is most common in the third or fourth decade. A terminal segment of esophagus 3 to 6 cm. long may narrow to a diameter of 1 or 1.5 cm.

As food lodges above the constriction, the tube gradually expands,

thickens, and lengthens, taking various forms with passage of time. The dilated sac is fusiform, then flask shaped, and finally sigmoidal (Fig. 1).


Cardiospasm must be distinguished from reflex spasm, benign stricture, peptic ulcer, congenital stenosis, and cancer.

The chief manifestations of cardiospasm are dysphagia, regurgitation, and retrosternal pain. During excitement or depression spasm is intense.

The lack of food causes anemia, weight loss, and eventually severe inanition.

Roentgen examination after a barium meal shows complete or incomplete blockade. The lower esophagus is narrowed and ends in a rounded cone, nipple, or stringlike process pointing to the left below the diaphragm.

Figure 1. Progressive development of esophageal sac



* Surgical treatment of cardiospasm. *Postgraduate Med.* 5:351-360, 1949.

CORRECTION OF ESOPHAGEAL DEFORMITY




Figure 2

SURGERY

An early stage of tension is sometimes relaxed by psychiatric care, antispasmodic drugs, particularly octyl-nitrite, and a bland, nutritious, non-residue diet. Established stricture is abolished by dilatation with a hydrostatic bag in 70% of cases and decreased in 20%.

Operation should be done when dilatation under direct vision is impractical or ineffective or when the diagnosis is uncertain and cancer suspected. Infants and children are not often helped by stretching technic but are usually greatly benefited by surgery.

A fluid diet is given for a few days before operation and the esophagus repeatedly rinsed with weak sodium bicarbonate solution. Shortly beforehand a tube is passed through the nose and gastric contents are aspirated. The tube is left in place until a day or two after surgical correction.

The abdomen is opened through a left epigastric incision, the stomach drawn down, and the left lateral hepatic ligament divided. The peritoneum over the esophagus is incised for 2 in. (Fig. 2a) and the esophagus mo-

bilized by blunt and sharp dissection (Fig. 2b). The left vagus nerve is drawn aside. To aid traction, a sling of gauze is placed around the esophagus (Fig. 2c), and 2 to 4 in. of the tube is pulled down into the peritoneal cavity.

A 4- or 5-in. cut is made through the muscular layers of the esophagus, continued over the cardiac region, and curved slightly upward toward the fundus (Fig. 2d). Longitudinal muscle fibers and the thin, somewhat adherent circular coat are cautiously divided until the underlying mucosa bulges outward (Fig. 2e).

To detect puncture of the mucosa, a fairly frequent but not serious accident, the body of the stomach is squeezed to force air and gastric juice into the esophagus. Any nick or tear is readily closed with silk.

Postoperatively the esophagus will usually contract to natural size, although after long deformity the sac may persist. A fluid diet is given for three or four days, then semisolid food. Before discharge from the hospital on the seventh or eighth day unrestricted meals are tried.

PERICARDIAL ASPIRATION is the most important initial treatment for cardiac tamponade due to wounds of the heart. Administration of blood and plasma is a desirable adjuvant. Facilities for open operation should be at hand, but with most wounds which cause cardiac tamponade and are not immediately fatal, aspiration alone will suffice. Mark M. Ravitch, M.D., and Alfred Blalock, M.D., of Johns Hopkins University, Baltimore, report 17 cases of stab or bullet wounds in the heart. Aspiration alone was adequate in 9 cases; 1 patient recovered after aspiration but later suddenly died. Two required operation; 5 died before aspiration could be done. The pericardial sac is pierced through the left fourth or fifth interspace parasternally by a 16- or 18-gauge needle.

Arch. Surg. 58:463-477, 1949.

Poison Ivy Management

J. B. HOWELL, M.D.*


Dallas

PROPER treatment of poison ivy dermatitis should give at least partial relief of itching, prevent secondary infection, and promote healing.

Itching is best controlled by topical therapy. The type of remedy chosen depends on the stage of the dermatitis. If the affected area of skin is moist

4-in. gauze bandage soaked in any one of several solutions prepared with distilled water (see table) for wrapping an arm or leg affected with poison ivy. Unstarched white sheeting or similar material may be substituted for the gauze bandaging if desired.

The wrapping is anchored between the fingers or toes (see illustration)



Method of applying wet dressing

with weeping blisters, a wet dressing should be applied. The eruption is not spread by the blister fluid, hence precautions against contact with unaffected skin are unnecessary.

J. B. Howell, M.D., employs 8-ply,

and wound so as to cover all the involved surface. Large bath towels are then tied over the gauze. The dressing should be kept sopping wet either by resoaking in the solution or by spraying with a syringe.

* Suggestions for management of acute poison ivy dermatitis. *Texas State J. Med.* 45:349-353, 1949.

DERMATOLOGY

If the face is involved, a mask is fashioned from several thicknesses of gauze mesh and applied wet to the head of the patient as pictured.

When the eruption involves extensive areas of the body, medicated baths (see table) kept at a temperature of 90° to 98° F. may be taken. The patient remains in the bath for fifteen minutes to several hours at a time.

If wet dressings or baths cannot be employed throughout the day, a lotion, emulsion, or tincture (see table) may be helpful if applied as often as the cycles of pruritus recur. The crusted lotion should be removed once daily by sopping the area carefully with a 2 to 4% solution of boric acid.

Ointments and stimulating medications are contraindicated. Preparations containing local anesthetics, such as nupercaine, surfacaine, butesin, or benzocaine, and thephorin are best avoided because of the risk of sensitization.

During the involuting, drying stage of the dermatitis, creams containing pyribenzamine, benadryl, or histadyl may be tried. Zinc oxide pastes with or without antipruritics are also beneficial after the weeping and exudation have subsided.

Drugs taken orally to alleviate itching are of little value. Aspirin or chloral hydrate may be of some benefit at bedtime. Opiates are not advisable.

If secondary infection does occur, the conventional therapeutic measures should be given. Poison ivy extracts, either parenterally or orally, are of no value in the acute stage of poison ivy dermatitis and may even be detrimental.

A few doses of 40 to 75 r units of

superficial unfiltered roentgen rays at five- to seven-day intervals may lessen itching, but usually not as efficiently as topical dermatologic remedies. Calcium gluconate, 10 cc.; strontium bromide, 1 ampule, intravenously every one to three days; or autohemotherapy, 5 to 15 cc., two to three times weekly may break the itching cycles and give slight relief.

SOLUTIONS FOR TOPICAL MEDICATION

Wet dressing solutions

1] Boric acid solution: Add 1 level tsp. boric acid crystals to every 2 glasses of water. Warm water dissolves the crystals more easily than cold water.

2] Burow's solution (1:20-1:30): Add 3 tsp. Burow's solution to every 2 glasses of water.

3] Potassium permanganate solution: Add 1 crushed potassium permanganate tablet (5 gr.) to every 2½ qt. water. Prepare fresh. The solution will stain clothing, tubs, basins, etc.; the stains can be removed from utensils with calcium oxalate, vinegar, or acetic acid. This solution is not suited for use about eyes or face.

4] Alibour water (modified):

	Gm. or cc.
Copper sulfate	0.6
Zinc sulfate	2.0
Camphor water, in sufficient quantity	
To make	100.0
(Dilute 1:16 to 1:25.)	

5] Thiersch solution (modified):

	Gm. or cc.
Boric acid	12.0
Salicylic acid	2.0
Menthol	2.0
Alcohol (188 proof)	50.0
Water, in sufficient quantity	
To make	1,000.0

- 6] Cornstarch solution: Add from 1 to 2 oz. cornstarch to each qt. water.
 7] Calgon solution: Add $\frac{1}{2}$ tsp. Calgon to every 2 qt. water.
 8] Tannic acid solution (2.5%).

Medicated baths

- 1] Cornstarch bath: Stir from $\frac{1}{2}$ to 1 lb. of refined, soluble cornstarch (preferably the unperfumed Linit starch) into a tub full of water.
 2] Potassium permanganate bath: Dissolve completely from 15 to 45 5-gr. potassium permanganate tablets in 1 qt. warm water. Use 1 qt. of the solution to each tub of water.
 3] Tar bath: Use approximately 3 to 4 tbs. Almay tar bath (a preparation of oil of cade in a sulfonated oil) to a tub of water. The patient may be painted with a tar preparation such as liquor carbonis detergens and placed in an ordinary bath, or the tar may be dissolved by stirring into the water 3 oz. of liquor carbonis detergens.
 4] Colloid bath: Cook 2 cups of bulk oatmeal in 1 qt. of water from 30 to 45 minutes in a double boiler; allow the mixture to cool for 15 minutes; add $\frac{1}{2}$ cup baking soda; pour the mixture into a gauze bag and tie the top; place in a tub of water at 90 to 96°F. The patient may stay in the tub 30 to 40 minutes expressing the oatmeal mash through the gauze and applying it over the body. The mash should be washed off thoroughly before the patient leaves the tub. Bulk oatmeal is preferable to the precooked type.

Lotions

- 1] Biborate lotion:
- | | Gm. or cc. |
|---|------------|
| Sodium biborate | 10.0 |
| Starch | 15.0 |
| Zinc oxide | 15.0 |
| Lime water, | |
| Rose water, each in sufficient quantity | |
| To make | 240.0 |
- 2] Calamine lotion (N. F. VI).

- 3] Lotion base:

	Gm. or cc.
Zinc oxide	15.0-20.0
Talc	15.0-20.0
Glycerine	10.0
Distilled water	80.0

- 4] Burow's lotion:

	Gm. or cc.
Burow's solution	15.0
Zinc oxide	30.0
Talc	30.0
Glycerine	24.0
Lime water	120.0

Emulsions (much improved by homogenization)

- 1] Bismuth emulsion:

	Gm. or cc.
Bismuth subnitrate	4.0
Zinc oxide	8.0
Lime water,	
Olive oil, each in sufficient quantity	
To make	240.0

- 2] Tragacanth lotion:

	Gm. or cc.
Tragacanth	2.0
Glycerin	0.33
Olive oil	60.0
Water, in sufficient quantity	
To make	240.0

- 3] Calamine liniment (N.F. or Pusey's modification).

Tincture

Glycerine	3%
Tannic acid	3-5%
Alcohol (95%),	
Distilled water, each in sufficient quantity	
To make	240.0 cc.

Pastes

- 1] Paste of zinc oxide (N.F.).
 2] Paste of zinc oxide to which one or more of the following may be added:
- | | |
|---------------------------|---------------------------------|
| Liquor carbonis detergens | 3-10% |
| Oil of cade | 1-3 % |
| Crude coal tar | $\frac{1}{2}$ -1 % |
| Ichthammol | 3-10% |
| Menthol | $\frac{1}{8}$ - $\frac{1}{2}$ % |
| Phenol | $\frac{1}{4}$ -1 % |
| Spirits of camphor..... | 2-3 % |

When Sunshine Is Harmful

J. M. HITCH, M.D.*

Raleigh, N.C.

THE benefits of sunshine are so greatly stressed that deleterious effects for some persons are frequently disregarded.

J. M. Hitch, M.D., warns that, besides the recognized hazards of sunburn, exposure to solar radiation may produce or aggravate skin diseases, especially with abnormal porphyrin metabolism or chronic actinic dermatoses.

Ultraviolet rays are responsible for most of the adverse reactions but a few are caused by heat from infrared rays. Some external chemical agents are photosensitizers.

EFFECTS OF ULTRAVIOLET EXPOSURE

Sunburn—Persons especially susceptible to sunburn should shield the skin by salol, quinine, or tannic or para-aminobenzoic acid in a greaseless base. Some commercial suntan lotions are valuable. Burned areas are soothed by oily or water-miscible emollients. To avoid toxic absorption antipruritic and analgesic drugs should be applied in very low concentration. Chills, fever, dehydration, and shock are combated by warmth, fluids, and sedation.

Photosensitivity—Among external factors that lower resistance to sunlight are coal-tar derivatives such as eosin, erythrosin, and some fluorescent products. Pyrethrum and oil of bergamot—a plant extract often used in perfume, cologne, and mosquito re-

pellants—may cause dermatitis if treated surfaces are exposed to sun.

The chief internal sensitizers are porphyrins in pathologic amounts due to hereditary metabolic defects, liver disease, or drugs, for example sulfonamides. However, the connection between skin disease and excess porphyrin is not always demonstrable.

Hydroa aestivale, or *summer eruption*, begins in the first four years of life and disappears at or soon after puberty. On cheeks, nose, and forehead a red papulovesicular eruption appears in successive crops as long as exposure continues. Itching and burning may be intense and scarring permanent.

Epidermolysis bullosa may be caused by ultraviolet stimulation in adult life. Vesicles and large bullae are seen on a red, hyperpigmented, hairy surface.

Eczema solare, first a papulovesicular inflammation, changes to weeping dermatitis and eventually a chronic verrucous form. Estrogens may help relieve symptoms of the disease in women with normal menstruation and in men.

From porphyrin-induced lesions the skin should be protected by ointments or clothing. Large oral doses of vitamin B complex and sodium para-aminobenzoate are given. Pyribenzamine may be administered systemically or applied over limited areas.

* The sun and the skin. North Carolina M. J. 10:299-304, 1949.

Xeroderma pigmentosum, due to a recessive gene, begins with deep freckling in early childhood. Atrophy, telangiectasis, keratosis, papillomatous growth, and carcinoma develop and are usually fatal before puberty. Sunlight should be avoided and roentgen or radium treatment applied to epitheliomas.

Chronic actinic dermatitis is rapid aging of skin by sunlight, wind, cold, heat, and ordinary senile changes. A patient in the second decade of life may appear old and weather-beaten. Irregular brownish pigmentation on neck, hands, and forearms alternates with patches of vitiligo, atrophy, wrinkling, dryness, telangiectasis, flat keratosis, and finally epithelioma.

Patients should be protected from the weather and keratoses removed by desiccation. All thickened, ulcerated, crusted, or tumorous areas should be biopsied. Routine semianual examination and treatment of

malignant growth may prolong life for many years.

EFFECTS OF INFRARED EXPOSURE

Miliaria rubra, or *prickly heat*, results from blockade of sweat glands by swollen squamous cells. Irritation is relieved by thin clothing, soda and starch baths, alcohol rubs, and local application of lotio nigra or lead and laudanum wash.

Urticaria and *angioneurotic edema* result from either heat or ultraviolet rays. The source is found by tests with a heating pad, infrared and ultraviolet lamps, and sunlight in cool air. Heat sensitivity may be overcome by increasingly warm baths. Antihistamines often give relief but in severe chronic cases sunlight must be avoided.

Erythema ab igne is reticulated erythema due to vascular dilatation by repeated exposure to heat. Diapedesis occurs and when hemosiderin is deposited the process is irreversible.

PRURITUS unchecked by other medication may be temporarily relieved by carbowax ointment containing 2% histadyl hydrochloride, a potent histamine antagonist. In 41 of 104 patients observed by Eugene S. Bereston, M.D., of Johns Hopkins University, Baltimore, itching subsided and some lesions healed as long as treatment was continued, but recurred later. Cases included instances of disseminated or localized neurodermatitis, contact or eczematoid eruptions, and irritation of anus, vulva, or scrotum. No toxic reactions developed.

J. Invest. Dermat. 12:157-158, 1949.

DERMATOPHYTOSIS may be treated effectively with an alcoholic solution of 2.5% 8-hydroxyquinoline. Such a preparation was used in treatment of 40 cases. Kurt A. Oster, M.D., and Milton J. Golden of Bridgeport, Conn., declare that results in light cases appear to be equal to those obtained with other antifungal substances and, in severe cases, superior.

Exper. Med. & Surg. 7:37-45, 1949.

Juvenile Cirrhosis of the Liver

PAUL D. KELLER, M.D., AND WILLIAM L. NUTE, JR., M.D.*

Washington University, St. Louis

CHILDREN may have hepatic cirrhosis at any age. Inherited defect or major derangement in some other organ system may be causative.

In the St. Louis Children's Hospital 40 cases proved by autopsy or operation were seen in twenty-six years.

Paul D. Keller, M.D., and William L. Nute, Jr., M.D., noted several major types. Forms not represented were syphilitic, pigmentary, lipoid, and zooparasitic.

Most cases of *obstructive biliary cirrhosis* result from congenital atresia of extrahepatic bile ducts. The stools are white or clay colored from birth, and jaundice soon appears. Infants usually remain surprisingly well nourished for weeks, but bleeding and portal obstruction eventually develop; the liver becomes palpable and spleen moderately large. Laboratory data show complete obstruction with little if any urinary urobilinogen.

If operation is neglected, death often occurs before the age of eight months. Rare factors are stone in the common duct, enlarged lymph nodes, viscid bile, and pressure from tumor. Hepatic tissues show interlobular fibrosis, bile stasis, and infiltration.

Nodular cirrhosis usually develops in children past infancy. Causes are obscure but possibly include infection, malnutrition, and sickle-cell anemia. In most of 11 cases reviewed,

Banti's syndrome was present, at times without anemia.

Most common manifestations are jaundice and gradual swelling of abdomen and legs. Children may have ascites, poor appetite, weakness, gradual weight loss, repeated infection, vague abdominal pain, and nosebleed. Symptoms vary and diagnosis depends on liver biopsy, which should be done by laparotomy. Tissues contain perilobular fibrosis, focal regeneration, and mononuclear infiltration without bile stasis.

Erythroblastosis fetalis may cause hepatomegaly and splenomegaly and varying degrees of cirrhosis. Many who recover probably have residual scars, but the 5 infants observed lived only a few days or weeks. Parenchymal injury of the liver, necrosis, biliary stasis, and hemosiderosis are found. Fibrosis is invariably diffuse.

Congestive cirrhosis in childhood generally follows a long period of intermittent cardiac decompensation and is overshadowed by the primary disease. In 1 case the fibrotic state was due to endophlebitis with partial block of the hepatic vein. The excess fibrous tissue is central to the lobule.

Toxic or postnecrotic cirrhosis is due to such agents as chloroform, salvarsan, carbon tetrachloride, and sulfonamide. The hepatic fibrosis in childhood is the same as in adult life. The liver becomes small and unevenly

* Cirrhosis of the liver in children. J. Pediat. 34:588-615, 1949.

nodular with bands of scar tissue. In the case observed, the liver damage was due to cinchophen contained in tablets eaten by the child.

Hepatolenticular degeneration is a curious hereditary disease affecting the liver and lenticular nuclei in the brain. A pigment ring may be found at the corneal limbus. Illness begins

in late childhood with gradual bulbar palsy, speech disorder, emotionalism, coarse tremors, rigidity, and contracture of hands, wrists, and feet. The mouth sags open and face appears idiotic long before the mind deteriorates to comparable degree. Liver changes are identical with ordinary portal cirrhosis.

Curare and Exercise for Poliomyelitis

W. D. PAUL, M.D., AND O. A. COUCH, JR., M.D.*

THE tight, painful muscles encountered in the acute phase of poliomyelitis are best treated by stretching exercises which utilize the full range of motion of the involved joints.

When spasticity is slight, these exercises may be carried out with little discomfort, but when the muscles are strongly contracted, tightness and pain usually prevent stretching sufficient to achieve good results. The use of curare permits satisfactory physical therapy.

W. D. Paul, M.D., and O. A. Couch, Jr., M.D., of the State University of Iowa, Iowa City, employ curare in the acute stages of poliomyelitis to relax muscles enough for adequate stretching. During the first day of therapy, 1.2 units of Intocostin per kilogram of body weight is given every eight hours, then 0.9 unit per kilogram every eight hours. Passive exercises are given by physical therapists thirty to forty-five minutes after each dose.

When curare is used, hyperextension must be avoided to prevent tearing the muscle fibers. Also, if the amounts and frequency of administration of curare are increased, respiratory distress is a possibility, especially when d-tubocurarine in oil is used.

Curare is used only to facilitate movement of the affected muscles and has itself no specific curative effect in poliomyelitis.

Curare and proper muscular exercises were employed for 24 patients with acute poliomyelitis. All were free of pain and muscle spasm within seventeen days. Another group of 21 patients with disease of similar severity was treated with the Kenny method of continuous hot pack routine without curare. Muscles could not be fully extended in most instances, and pain and tightness often persisted as long as five months.

* Preliminary report on the treatment of anterior poliomyelitis with exercise and curare. Arch. Phys. Med. 30:277-285, 1949.

Heart Disease and Pregnancy

JOHN J. SAMPSON, M.D.*

University of California, San Francisco

WOMEN with heart disease are bearing children more safely than ever before. Among 119 deliveries in a university clinic in the past twelve years, not one mother or baby died.

In all cases the paramount consideration is maternal risk of death or of permanent invalidism.

Hysterotomy is seldom warranted and, after the third lunar month, gestation is not often terminated by operation. After the eighth month, induced labor is rarely necessary as the circulatory burden then does not increase appreciably.

Unfavorable circumstances are active rheumatic fever within six months before conception, arrhythmia indicative of advanced cardiac disease, and age over thirty-five years.

Valvular lesions are relatively unimportant unless circulation is mechanically obstructed.

Cardiac dysfunction of Class I or II is relatively unimportant, but mortality rates almost triple with Classes III and IV. Hypertension involves no risk unless toxemia develops. Heart failure during pregnancy, unless due to a transient cause such as pneumonia, is an interdiction to subsequent childbearing.

Ordinary labor causes no particular strain, but when Class III or IV impairment already prevails a sudden disturbance during delivery may easily

precipitate heart failure; if such an event is known to have occurred, cesarean section may well be the advisable procedure.

Reasons for and against continued gestation must be balanced with care in each case, explains John J. Sampson, M.D.

For instance, a woman in the second month of pregnancy had Class III cardiac involvement. She had been more dyspneic since birth of the first baby and probably would not be able to care for two children. Rheumatic heart disease, which started thirteen years previously, had recently recurred.

But a second child was much desired, her religious beliefs opposed abortion, and the rheumatic exacerbation had subsided. She was certain of good medical supervision and household help before and after delivery. Pregnancy was allowed to continue to term, the heart did not fail, and no complications were encountered.

During pregnancy complete bed rest, digitalis, diuretics, and a low-salt diet may be required for women with heart failure. Quinidine is given to prevent or correct arrhythmia; antibiotics are used for infection. Oxygen may be needed, especially during labor.

Nitrous oxide is never given for deep anesthesia.

Bed rest should be continued post

* Heart disease as a complication of pregnancy. *California Med.* 70:383-390, 1949.

partum longer than usual. Oxytocic drugs may raise venous pressure dangerously and are not used.

As the circulation is likely to be overloaded by postpartum return of water to the circulatory system with consequent transient increase in blood volume, a mercurial diuretic or blood-letting measures may sometimes be necessary.

Any deviation from the natural circulatory changes of pregnancy may show imminent heart failure. Vital capacity, which should be measured often, commonly increases by about 10 to 15%.

Venous pressure in the arms rises only 1 or 2 cm. of water after the twenty-fifth week but increases in the legs for the last thirty weeks.

Circulation time diminishes from the twentieth to the thirtieth week. Pulse pressure is elevated in the last four months; blood pressure falls slightly in the second trimester and recovers in the third.

During gestation blood volume increases and hemoglobin percentage consequently drops. Immediately after delivery blood volume shrinks. Hemoconcentration is recovered in the next three to five days.

RELIEF FROM DYSMENORRHEA can often be obtained with magnesium gluconate. W. J. Rawlings, M.D., of Melbourne found that oral administration alleviated premenstrual distress for 17 of 19 patients, and menstrual pain for 14 of 18. Magnesium gluconate is given in doses of 1.3 gm. daily. For premenstrual distress, treatment is started seven days before onset of the menses and continued through the first day of menstruation. Treatment for menstrual distress is begun four days before onset and continued for the first three days of the period. If relief is not prompt, two or even three doses a day are prescribed.

M. J. Australia 1:61-64, 1949.

BASAL METABOLISM DURING MENOPAUSE decreases because of lessened thyroid activity, asserts Mary E. Collett, Ph.D., Western Reserve University, Cleveland. Small doses of estrin raise the BMR slightly and reduce hot flashes, but large doses depress the BMR and increase hot flashes. Basal metabolic rates covering 30 cycles were determined for one subject over a thirteen-year period together with eighteen tests at age sixty. At age thirty-five the cyclic BMR pattern showed two low points marking the menstrual period and ovulation, and two high areas presumably indicating periods of estrin and progesterin secretion. The pattern was the same at forty-five but the progesterin period was shorter. At forty-six and forty-seven, estrin and progesterin periods became irregular and finally the progesterin period disappeared.

J. Applied Physiology 1:629-636, 1949.

Medical Forum

Discussion of articles published in MODERN MEDICINE is always welcome. Address all communications to The Editors of MODERN MEDICINE, 84 South 10th St., Minneapolis 3, Minn.

Basket for Removing Ureteral Stones*

TO THE EDITORS: I agree with Dr. George R. Livermore's condemnation of using metal instruments for removing ureteral stones. Dr. Livermore's comment appeared in his discussion of Dr. W. P. Morton's basket for that purpose (Apr. 15, 1949, p. 76).

To insert a stone catcher into an otherwise functioning ureter and then drag a concretion through the narrow passage lined with delicate epithelium is no more sensible than the anal insertion of a grasping forceps to remove ordinary rectal residues.

New risks are added each time someone has glamorized another gadget and an instrument maker has been found to promote its sale. These fancy proposals have not even brought a satisfactory method for dilating the ureter to permit easy insertion of the gadgets intended to bring out a stone.

Ordinarily, the ureter permits passage of calculi small enough to enter the channel if proper dilatation by cystoscopic approach from below is provided. Urologists agree on this. However, the instrumentarium and methods of dilating are usually so complex that it is impossible to avoid undue pain, trauma, and infection.

*MODERN MEDICINE, Jan. 15, 1949, p. 71.

That being the case, it may be permissible to resurrect a cystoscopic procedure, improvements on the so-called Braasch type of direct water vision cystoscope.

Back in July 1928, the *Journal of Urology* published my article on "Ureteral Dilating Bougies and Cystoscope," which describes the simplest and, to this date, safest instruments for dilating the male ureter to facilitate passage of a calculus. The cystoscopic sheath is of the direct water vision type with a large oval lumen, unobstructed inside by any ridges. The catheterizing attachment extends the lumen of the sheath proximally by about 36 mm. It has a large cut-out on one side covered by an encircling rubber dam, secured against leakage by an external compression sheath. A perforation allows insertion of the bougie or catheter directly into the lumen of the cystoscope. The proximal end is closed with a convenient bubble draining ocular which permits visual control.

The bougies are of the Garceau type, but larger, and have shorter, stouter conical tips, tapering quickly to full size at about 6 or 7 cm., then continuing the full length in sizes of 12 to 18F. The tips of the bougies are well rounded but not bulbous, so that they are not likely to get caught in a fold of the ureteral mucosa, as often

happens with more pointed catheters. Consequently, the tip usually finds its way around a calculus where an ordinary catheter would be blocked and provides added dilatation where it will do the most good. Immediate drainage with relief of the renal colic usually results.

The patient remains ambulatory. Time is allowed for the calculus to drop down and be expelled into the bladder.

If the obstruction recurs, the dilatation may be repeated. If after an interval of ten to fourteen days, the calculus has not been expelled, dilatation is done with a larger bougie. The tip of the bougie may previously be dipped into melted beeswax to obtain scratch marks indicating the presence of the calculus. Bougies sizes 16 and 18F should follow if necessary but only at intervals of twelve to fourteen days in order to give the ureter a chance to recuperate. Common sense must be used, depending on how readily the ureter yields to dilatation. By introducing only one bougie at a session, trauma and infection are kept at a minimum.

This type of direct water vision cystoscope is better tolerated and less painful in the neck of the bladder than the ordinary catheterizing cystoscope with indirect telescope. Deep sedation and anesthesia are rarely needed. The large, straight bougie is the most effective ureteral dilator. It cannot break or get caught. It never snags the tissues. You can always pull it out intact—not so with bulbs and mechanical dilators.

The catheterizing attachment is really the key to the whole procedure, which makes it possible to get an 18F

bougie into the ureter through a 26F cystoscope, though the 28F sheath makes it easier. The attachment is so simple in construction that I have always made that part myself.

The bougies were made specially by Porges in Paris. If someone can persuade our American ureteral catheter manufacturers to make similar bougies, I will be glad to provide details. As it is, the largest ureteral catheter can be inserted through this cystoscope with greater facility and with less discomfort to the patient than with any other cystoscope. The American Garceau catheters, 12 and 14F may be used, preferably with a suitable stylet, to start the dilatations. Ultimately sizes 16 and 18F are essential when the stone is large and slow in passing. For permanent results in dilating ureteral strictures, an 18F bougie is necessary.

I am not trying to sell cystoscopes but, like Dr. Livermore, would like to help urologists do better and safer cystoscopies. The technic can be standardized. The demand for making the instruments must come from the profession.

RAYMOND L. SCHULZ, M.D.

Los Angeles



"Hmmm."

Diagnostix

Here are diagnostic challenges presented as they confront the consultant from the first clue to the pathologic report. Diagnosis from the Clue requires unusual acumen and luck; from Part II, perspicacity; from Part III, discernment.

Case MM-148

THE CLUE

ATTENDING M.D.: I would like you to see a seventy-year-old man who has been losing weight and feeling tired for seven months. Three months ago he had nausea and abdominal pain and since then diffuse, indefinite abdominal discomfort has persisted. Three weeks ago jaundice appeared with no colic and within the past few days he has vomited a small amount of blood. There is a palpable mass in the upper abdomen.

VISITING M.D.: What is the nature of this mass?

ATTENDING M.D.: I believe it is the liver. The left lobe of the liver is harder than the right and x-ray studies of the gastrointestinal tract, while not revealing intrinsic abnormality, show a mass compressing the stomach and duodenum.

VISITING M.D.: Are there other pertinent facts in the history?

above the umbilicus and left lobe is distinctly outlined. There is some ascitic fluid.

VISITING M.D.: Of course at this age, with progressive illness of about six months and a mass, one thinks of cancer. I do not believe it is carcinoma of the head of the pancreas. It could be a primary hepatoma. Is the patient alcoholic or does he have cirrhosis of the liver?

ATTENDING M.D.: We have no evidence of cirrhosis of the liver. For a number of years the patient has drunk whiskey, but not excessive quantities, 4 or 5 highballs a day at the most, and usually only 1 or 2.

VISITING M.D.: One must, of course, consider an amebic infection of the liver, with abscess and necrosis. Yet, that does not seem likely.

ATTENDING M.D.: We know he has gallstones, but the story is not that of painful obstruction due to common duct stone.

VISITING M.D.: What are the laboratory findings?

PART II

ATTENDING M.D.: There was a positive guaiac reaction of one stool specimen. The patient has become more fatigued and has lost 40 lb. Roentgenograms of the abdomen reveal several gallstones; the liver edge just



PART III

ATTENDING M.D.: The total proteins are 6 gm. per 100 cc. with 3 gm. of globulin and 3 gm. albumin. Alkaline phosphatase is 19 Bodansky units per 100 cc. Cephalin floccula-

(Continued on page 78)



**For
mixed
infections**

Over 500 cases of impetigo treated with Furacin have now been reported in the literature. Several investigators report good results in over 90% of their cases, often within an average of seven days. Of 30 cases of ecthyma reported, good results were obtained in 24 within the average time of eight to ten days. Sensitization averaged under 5 per cent. Furacin® brand of nitrofurazone is available as Furacin Soluble Dressing (N.N.R.) and Furacin Solution (N.N.R.) containing Furacin 0.2%. These preparations are indicated for topical application in the prophylaxis or treatment of infections of wounds, second and third degree burns, cutaneous ulcers, pyodermas and skin grafts. *Literature on request.*

EATON LABORATORIES, INC., NORWICH, N. Y.

Dillane, W. B. et al.: *Treat. Serv. Bull.* 2:47, 1947 • Downing, J. G. et al.: *J. A. M. A.* 133:299, 1947 • Downing, J. G.: *Am. Pract.* 2:357, 1948 • Downing, J. G. et al.: *New England J. Med.* 239:62, 1948 • Eichenlaub, F. J. et al.: *M. Ann. District of Columbia* 17:452, 1948 • Johnson, H. M.: *Arch. Dermat. & Syph.* 57:348, 1948 • Miller, J. et al.: *New York State J. Med.* 47:2316, 1947 • Robinson, H. M. et al.: *South. M. J.* 40:409, 1947.

Edocet
Experientia



Picture the patient's progress

...with photograph...after photograph


Easy enough to show them, too. Just load the new Kodaslide Table Viewer with 35-millimeter or Bantam slides of cardboard or double-glass, or a mixture of both. Plug in—AC or DC. Then . . . on with the showing. This ingenious combination of projector, slide changer, and screen gives brilliant 4.8X enlargement, even in a fully lighted room. Use it to check slides . . . for consultation with colleagues . . . in discussion with patients or students . . . during case studies. For further information, see your nearest photographic dealer . . . or write to Eastman Kodak Company, *Medical Division*, Rochester 4, N. Y.

Major Kodak products for the medical profession



X-ray films; x-ray intensifying screens; x-ray processing chemicals; electrocardiographic papers and film; cameras—still- and motion-picture; projectors—still- and motion-picture; enlargers and printers; photographic films—color and black-and-white (including infrared); photographic papers; photographic processing chemicals; synthetic organic chemicals; Recordak products.

"Kodak" is a trade-mark

Serving medical progress throu



*Two Kodak Vari-Beam
Standlights are arranged at
this angle, equidistant
from the operative field.*



Photography and Radiography

Kodak

WASHINGTON LETTER

tion is 3 plus. Prothrombin time is 22 seconds, normal is 18. The x-ray examination of the esophagus did not reveal varices.

VISITING M.D: What has been the color of the stools?

ATTENDING M.D: Some stools were quite pale; others apparently contained pigments.

VISITING M.D: (*Examining patient*) I cannot feel the spleen. The prostate is normal in size. I believe the patient has obstructive jaundice probably due to cancer.

ATTENDING M.D: The hemoglobin is 14 gm. per 100 cc.; white blood cell count is 15,000; urine contains bile. Serum cholesterol is 200 mg. and the nonprotein nitrogen 35 mg.

VISITING M.D: This does not suggest extensive liver damage. I believe the patient has a hepatoma. I would suggest a needle biopsy of the liver.

PART IV

ATTENDING M.D: Biopsy reveals hepatic cirrhosis of alcoholic type and hepatoma.

Washington Letter

(Continued from page 44)

a constitutional majority, a majority of all members. That would be 49 senators or 218 representatives. However, the plan dies automatically if Congress adjourns before August 19.

If the plan goes through, it will mean settlement by presidential edict of an issue on which Congress itself couldn't agree. The creation of a department of welfare has been proposed at the last three congresses.


Movie on Breast Cancer

National Cancer Institute and American Cancer Society are producing a movie on breast cancer, intended for women's clubs and organizations. It is a color film and will include a simple technic for periodic self-inspection of the breasts. Also to be released for distribution in the fall is a professional film of thirty minutes depicting technics of breast cancer diagnosis. . . . Studies at the Cancer

Institute indicate that eclampsia, which accounts for one-fourth of all maternal deaths in this country, has some relation to hormonal or endocrine imbalance. Dr. Alexander Symeonidis is conducting experiments on the subject. . . . The institute has granted \$8,500 to the California Public Health Department to study the factors in the human environment that may influence the development of lung cancer. The employment records of 500 lung cancer cases will be studied and, when a possible occupational exposure is indicated, continuing investigations will be made. This was one of 36 special cancer control grants totaling more than half a million dollars. . . . Seventy-four of the country's 79 medical schools and 36 of the 40 dental schools are receiving grants in cancer training programs. The grants, of \$5,000 to \$25,000, may be renewed annually.

You Can Forget About Sterilization

Ritter's SYNCHRONOUS TIMER Sterilizes Automatically!



● Now you can concentrate on your important income-producing work. This amazing synchronous timer makes your Ritter sterilizer a faithful, completely automatic servant. The timer operates *only* when water reaches the boiling point. You know positively that your instruments have been fully sterilized for the required time. Why bother with sterilization details? Let a new Ritter DeLuxe Hydro-matic Sterilizer take care of them for you, *automatically*.

MORE EXCLUSIVE RITTER FEATURES

- Automatic Water Supply
- Automatic Safety Switch
- Automatic Water Level
- Automatic Water Sterilization

Write for Catalog. Describes 6 types with many new features.

**Model "E-3" 16" New Timer
Assures Complete Sterilization**

FOR ADVANCED EQUIPMENT
LOOK TO

Ritter
COMPANY INCORPORATED
RITTER PARK, ROCHESTER 3, N. Y.



If vacationing in New York State, visit our plant and see quality Ritter equipment manufactured.



As the Incidence of Scabies Rises

The virtues of Kwell Ointment are discussed in these articles:

1. Wooldridge, W. E.: The Gamma Isomer of Hexachlorocyclohexane in the Treatment of Scabies, *J. Invest. Dermat.* 10:363 (May) 1948.
2. Nieldman, M. L.: Treatment of Common Skin Diseases in Infants and Children, *J. Pediat.* 32:566 (May) 1948.
3. Cannon, A. B., and McRae, M. E.: Treatment of Scabies, *J.A.M.A.* 138:557 (Oct. 23) 1948.
4. Goldman, L., and Feldman, N. D.: Human Infestation with Scabies of Monkeys, *Arch. Dermat. & Syph.* 59:175 (Feb.) 1949.
5. Fox, E. C., and Shields, T. L.: Résumé of Skin Diseases Most Commonly Seen in General Practice, *J.A.M.A.* 140:763 (July 2) 1949.

The specific, positive scabicial properties of Kwell Ointment are employed to particular advantage during the warm summer months with their concurrent increase in the incidence of scabies. Kwell Ointment usually eradicates scabies infestation with a single application. Its action is prompt, and secondary dermatitis or skin irritation rarely follows its use. Its extreme blandness makes its application permissible even in the presence of secondary infection.

Kwell Ointment owes its scabicial action to the gamma isomer of benzene hexachloride—0.5% in a vanishing cream base. This parasiticide is specifically lethal for *Sarcoptes scabiei* yet in the concentrations employed it is nontoxic to man. Available on prescription at all pharmacies in 2 oz. and 1 lb. jars.

C.S.C. Pharmaceuticals

A DIVISION OF

COMMERCIAL SOLVENTS CORPORATION • 17 E. 42nd St., N. Y. 17, N. Y.

KWELL[®] OINTMENT



Short Reports

SURGERY

Shunting Effect of Trauma on Gastric Circulation

Circulation of blood through the vessels of an ulcerated stomach is modified by surgical operation. When the peritoneal cavity is opened and the stomach is handled, arteriovenous anastomoses open in the region of the submucous plexus of vessels and stop active circulation through the small vessels of the mucous membrane. The shunting effect of trauma was demonstrated by Drs. A. E. Barclay and F. H. Bentley of the Nuffield Institute for Medical Research, Oxford, England. Arteries of pieces of stomach removed at operation and of cadaveric stomachs were injected with colloidal silver iodide. Cadaveric specimens revealed a rich vascular network of fine arterioles and capillaries running perpendicularly through the gastric mucosa toward the surface; these vessels were not filled in the operatively removed specimens. At 6 operations surface veins were observed from the moment the peritoneal cavities were opened. Within two or three minutes the veins coursing along the anterior wall of the stomach changed from purple-blue to dusky pink. Oxygen saturation of blood before and after the color change was 74% and 91%, respectively, confirming a belief that the blood passed directly from the arterial to venous side through an arteriovenous shunt. The anastomoses are probably under control of the autonomic nervous system, for

when the sympathetic outflow to the stomach was blocked by spinal anesthesia the shunts did not open and the mucosal vessels were completely filled.

Gastroenterology 12:177-183, 1949.

EDUCATION

Austrian Physicians Idle

Because of overcrowding in the medical profession Austrian high school students have been asked by the Association of Physicians not to enter medicine. Some 1,500 doctors are said to be unemployed at present.

PHYSICAL MEDICINE

Intraarticular Temperature as Measure of Joint Reaction


Accurate evaluation of the degree of synovial inflammation with rheumatoid arthritis is facilitated by determination of the intraarticular temperature. Drs. Steven M. Horvath and Joseph L. Hollander of the University of Pennsylvania, Philadelphia, assert that intraarticular temperatures correlate with clinical activity of arthritis more closely than surface temperatures over the joint. With degenerative joint diseases, intraarticular temperatures are higher than would be anticipated from the relative arthritic activity. Temperature within the joint falls when hot packs are applied over the joint and rises with application of cold packs. These responses are greater in winter than in summer.

J. Clin. Investigation 28:469-473, 1949.

Harvey Cushing's outstanding contributions
in the field of endocrinology and their
application to the study of medicine
are well recognized. The Armour Laboratories,
as a pioneer in the field of endocrinology,
is deeply appreciative of Dr. Cushing's
fundamental researches in endocrinology.

*Tenth in the series, PORTRAITS OF PIONEERS in
Endocrinology. A full-color reproduction of this painting
suitable for framing, is available upon request. On your
professional letterhead, please address:*

 **ARMOUR**
Laboratories
CHICAGO 9, ILLINOIS



Harvey W. Cushing, M.D. 1869 - 1939

Destined to greatness by his intellect, character and energy, Harvey Cushing gave much to his fellow men and in return he received tremendous success and acclaim. A great scientist, he nevertheless was a master of the art of clinical medicine. Although pre-eminently known as a neuro-surgeon, he made fundamental researches in the functions of the endocrine glands. Particularly outstanding were his investigations of the secretions of the anterior pituitary lobe and their clinical effects.

Harvey Cushing was born in Cleveland, April 16, 1869. He graduated from Yale College in 1891 and from Harvard Medical School in 1895. He interned at Massachusetts General Hospital and

then became resident in surgery under the famed Doctor Halsted at the newly opened Johns Hopkins Hospital. Early in his career he determined to enter the then infant field of neurological surgery. He went abroad and studied under Horsley, Kocher and Sherrington. On his return he did experimental work at Johns Hopkins and published his epochal book, "The Pituitary Gland and Its Disorders". In 1912 he was made Professor of Surgery at Harvard and head of the surgical service at Peter Bent Brigham Hospital. Here he worked prodigiously for many years developing the field of neurological surgery, investigating endocrine function, and writing pertinent articles and monographs.

SHORT REPORTS

ORTHOPEDICS

Osteogenesis Produced by Extract of Bone

A chemical substance, when injected into skeletal muscle, may stimulate bone formation. Such a substance is contained in alcoholic extracts of the growing ends of postfetal bone, assert Dr. Joe Hartley and associates of Mount Sinai Hospital, New York City. Extract prepared from the ends of long bones removed from rabbits less than five weeks old was injected into 11 rabbits. Microscopic examination fifty-three days later revealed new bone formation in 3.

J. Mt. Sinai Hosp. 15:383-387, 1949.

PSYCHIATRY

Allergy in Psychotic Reactions

The mentally ill are less susceptible to hay fever than the sound in mind. Incidence of major allergy among psychotic patients is much less than among individuals free from psychiatric disorders. At the height of the ragweed season 1,875 patients and 757 healthy employees of the Veterans Administration Mental Hospital, Coatesville, Pa., were examined for symptoms and signs of major allergy. Positive physical findings of allergic response were found in 13% of the employees, report Drs. Robert M. McAllister and Arthur O. Hecker of the hospital staff. Incidence among schizophrenic patients was 2.9%; among manic-depressive patients, 1.4%; and among patients with organic psychoses, 3%. Incidence of allergy among epileptics was 13%, the same as among the healthy employees.

Am. J. Psychiat. 105:843-848, 1949.

METABOLISM

Hormonal Stimulus for Growth

Biologic effectiveness of endogenous steroid hormones may be suppressed by several folic acid antagonists. Drs. Roy Hertz and William W. Tullner of the National Cancer Institute, Bethesda, Md., report that a purine analogue, 2, 6-diaminopurine, inhibits estrogen-induced growth in the genital tract of the female chick. The inhibition is largely reversible by adenine, known chemically as 6-aminopurine. However, as much as 5 mg. of folic acid does not reverse the inhibition produced by 10 mg. of 2, 6-diaminopurine. The degree of response obtained in a tissue which is under maximal hormonal stimulus for growth can be quantitatively determined by folic acid and adenine and their respective inhibitory analogues.


Science 109:539, 1949.

ENDOCRINOLOGY

Adrenals and Glycogenesis

Conversion of protein to glycogen is effected by mediation of the adrenal glands. Glycogenesis is impaired in event of adrenal cortical insufficiency. This metabolic defect may be corrected, suggest Dr. Jorge Awapara and associates of the University of Texas, Houston, by administration of 17-hydroxydehydrocorticosterone (Compound E). Liver glycogen is increased when adrenalectomized rats are given Compound E. The dicarboxylic amino acids are decreased and alanine is increased. Changes in amino acid concentration and distribution in muscle are not impressive.

Endocrinology 44:378-383, 1949.



When complaints of bloating and belching suggest dyspepsia caused by biliary insufficiency, the choleretic-digestant-laxative action of ZILATONE Tablets usually provides prompt relief.

ZILATONE Tablets combine bile salts compound with extract of cascara sagrada, phenolphthalein, nux vomica, pepsin, pancreatin, and capsicum. They improve the quantitative and qualitative supply of bile, aid fat metabolism, assist carbohydrate digestion, and stimulate intestinal motility and bowel evacuation.

zilatone TABLETS

Available at all pharmacies in packages of 20, 40 and 80 tablets.
Samples and literature on request.



DREW PHARMACAL CO., INC., 170 VARICK STREET, NEW YORK 13, N. Y.

SHORT REPORTS

NEUROSURGERY

Uptake of P^{32} by Cancer of the Brain

Avidity of malignant brain tumors for circulating inorganic phosphate may make possible significant intracellular concentrations of radioactive phosphorus. Dr. Theodore C. Erickson and associates of the University of Wisconsin, Madison, gave intravenous inorganic phosphate containing P^{32} to 10 patients with malignant intracranial tumors. Some biopsy specimens had nineteen times as great concentration of radioactivity in tumor tissue as in healthy brain tissue. Chemical fractionation of tissue phosphorus revealed a small but significant concentration in the nucleoprotein of both cytoplasm and nucleus.

J. Lab. & Clin. Med. 34:587-591, 1949.

PUBLIC HEALTH

WHO Membership

Ecuador and Costa Rica have recently joined the World Health Organization, which now has a membership of sixty.

GENETICS

Predetermination of Sex

The time factor between insemination and ovulation apparently directly influences the sex of offspring. The sex ratio is presumed to be approximately equal in the middle of the fertility period; female preponderance is probable in the early stages, male in the later. Studies in rats reveal that delaying insemination until varying hours after the expected time of ovulation, increases the ratio of male to female offspring from 100:1:100 to between 149:100 and 225:100. Drs. Deryl Hart and James D. Moody of Duke University, Durham, N.C., believe that the same factor operates in determining the sex of human beings. An analysis of more than 65,000 pairs of dizygotic twins reveals in each year's group approximately 27% more like-sexed than unlike-sexed pairs. This constant surplus is inconsistent with chance alone and probably results from inseminations early and late in the fertility period.

Ann. Surg. 129:550-571, 1949.

EXPERIMENTAL SURGERY

Occlusion of Large Arteries

After aneurysmal dilatation, occlusion of a large artery by metallic band is often unsuccessful because atrophy and erosion of the vessel wall under the band are provoked. The reaction of tissues to polythene has been used by Dr. F. W. Cooper, Jr., and associates of Emory University, Ga., in conjunction with a tantalum band in operations on the aortas of dogs. The fibrosis induced by the plastic consolidates constriction of the band.

Surgery 25:184-190, 1949.



New



STERILE

CONVENIENT

ABSORBENT

UNIFORM

ECONOMICAL

ACTUAL SIZE 1 1/4" DIAM.

RED CROSS COTTON BALLS IN A NEW CONVENIENT FORM

FOR applying medication, cleansing wounds, parenteral therapy, pressure dressings, preoperative skin preparation, prehypodermic injection.

COMPACT CARTON OF 65 ready for instant use, convenient in your bag or on office shelf.

RED CROSS COTTON BALLS

What Can Be Done for the Hay Fever Patient?

(Continued from page 50)

tadyl and 0.5% pyribenzamine nasal solutions. The latter is now furnished in a nebulizer and considered less irritating. Preparations that contain ephedrine or epinephrine should not be used.

The patient must realize that instillation treatment is allowed for only a short period of time. Nothing is worse than a vasomotor rhinitis medicamentosa.

Sedatives are also of value in some cases, but many clinicians don't give them a thought now that so-called antihistamines have been placed on the market.

Which antihistamine is best suited for the patient? Will there be any side reactions? These questions are difficult to answer.

ANTIHISTAMINE THERAPY

Many publications have clearly revealed the history, chemistry, and laboratory reactions of the antihistamines. Numerous clinical papers have also been published, but unfortunately they praise the antihistamines altogether too much. It would be better if the physician could obtain from a few very thoroughly controlled studies the realization that the antihistamines should be offered with many words of caution. Common side effects are shown in Table 1.

A recent survey by the Research Department of Modern Medicine Publications has brought some interesting facts to light. The report, based on information received from

720 doctors distributed throughout the United States, reveals that treatment of the patient with hay fever rests chiefly in the hands of the general practitioner, but involves all specialists. The survey shows that thirteen different antihistaminic drugs were prescribed by the doctors questioned.

Most of the doctors used those antihistamines which were first introduced to the medical profession. There is, however, no reason why all the preparations now available should not be considered in therapy. The confusion which exists in the minds of many clinicians might then be cleared up. Some doctors feel that it is best to limit themselves to prescribing a few drugs, letting the results be what they may. Others have tried the method of rotation. This system has given better help than might be expected, but patients have sometimes thought that the doctor did not know much about what he was doing when he prescribed one antihistamine after another!

Now, however, a mass of information is pouring in to indicate that a certain pattern of approach can be followed in giving antihistamines to the hay fever sufferer. The more one studies the clinical behavior of these drugs the more one learns that they can be put to good use. The various antihistamines are listed, according to groups, in Table 2.

Group 1—The derivatives of ethylenediamine may be tried first. They

Detailed Dosage Data For Physicians

A 20-page brochure prepared exclusively for the medical profession presents busy physicians with detailed dosage information relating to quantity, frequency and duration of administration in relation to menses, as well as indications, rationale, etc., regarding ERGOAPIOL (Smith) with SAVIN. This time-tested uterine tonic is thoroughly described in this brochure, "Menstrual Disorders—Their Significance and Symptomatic Treatment." A copy, available to physicians only, will be supplied on request. Ethical since its inception, ERGOAPIOL (Smith) with SAVIN is dispensed only on your prescription.

INDICATIONS: Amenorrhea, Dysmenorrhea, Menorrhagia, Metrorrhagia, and to aid involution of the postpartum uterus.

GENERAL DOSAGE: 1 to 2 capsules, 3 to 4 times daily—as indications warrant.


In ethical packages of 20 capsules each, bearing no directions.



Ethical protective mark, M.H.S., visible only when capsule is cut in half at seam.

ERGOAPIOL (SMITH) **WITH SAVIN**

Martin H. Smith Co. • 150 Lafayette St. • New York 13, N.Y.



include neohetramine for rather mild symptoms, neo-antergan for moderate distress, and pyribenzamine for more severe manifestations. These antihistamines may fail to give satisfactory results, although they are relatively unlikely to cause side reactions.


Group 2—Drugs with a “thenyl ring” may then be substituted. They are thenylene, histadyl, diatrin, tagathen, and chlorothen. The chemical structure of the latter two is similar, but tagathen comes in the form of a coated tablet and the other does not. Some investigators feel that this makes a difference in the efficacy of the preparations. Studies to explore this possibility are in progress.

Mild to moderate sedation may accompany the administration of the antihistamines of this group, but often the side effects are actually of benefit in control of the nervous hay fever patient.

Group 3—Miscellaneous drugs are trimeton, pyrrolazote, and thephorin.

The action of trimeton and pyrrolazote is much the same, although the former may be a little less effective and the latter produces some sleepiness. Thephorin stands alone. It gives satisfactory results in many cases but has a stimulating effect and, when taken toward the end of the day, may cause insomnia. However, the tired hay fever sufferer may respond well to thephorin during the day.

Group 4—Finally, there are the ethanolamine derivatives, namely, decapryn and benadryl with its modification called hydryllin. All are good antihistamines but they yield a rather high incidence of drowsiness. This feature is no contraindication to their use, however, since many individuals with hay fever are highly irritable and



“Let's face it Chief, you're allergic to feathers!”

a
therapeutic
key
in
arthritis

ERTRON[®]

steroid
complex,
Whittier

Administration of Ertron, Steroid Complex, Whittier often provides a therapeutic key to the difficult problem of arthritis. Controlled medication with Ertron, accompanied by proper adjunctive measures, generally fosters an improvement in joint function over a wider range of movement and a reduction in local swelling, stiffness and pain. Aspects of systemic improvement are evidenced in increased appetite, better nutrition and a gain in weight.

These arthrokinetic and anti-arthritic effects of Ertron stem from the Whittier process—electrical activation of heat vaporized ergosterol. Clinical improvement has been shown in more than 80 per cent of cases treated with Ertron during thirteen years of investigation.

Ertron is supplied in bottles of 50, 100 and 500 capsules, and Ertron Parenteral in packages of six 1 cc. ampuls. Each capsule contains 5 milligrams of activation-products having antirachitic activity of fifty thousand U.S.P. units. Each ampul contains activation-products having antirachitic activity of five hundred thousand U.S.P. units, in sesame oil. Biologically standardized.



Whittier

LABORATORIES

Division Nutrition Research Laboratories, Chicago 30, Illinois



SPECIAL ARTICLE

require definite sedation. Some patients, therefore, respond well to the drugs as daytime medication; others need them only at bedtime and during the night.

The antihistamines of Group 1 may be employed during the day, starting with small doses after meals. If side reactions are insignificant, the drugs can be administered before meals, often with better effect.

Drugs from Group 2 may be used both day and night.

A combination can be worked out with the antihistamines of Group 3. Thephorin is taken early in the day, and trimeton or pyrrolazote at the end.

The antihistamines of Group 4 are frequently efficacious toward the end of the day because they promote sleep.

No matter how good the effects from any one preparation, it must be remembered that complications cannot be avoided. Skin eruptions resembling atopic dermatitis have appeared. The many cases of resulting asthma have led to the great emphasis now being placed on specific pollen therapy.

POLLEN THERAPY

This form of treatment does not consist of the simple administration of a pollen extract to all hay fever patients but includes a careful study of each case to determine the cause of the disease and the best remedy.

Too many physicians fail to acquaint themselves with the pollens of their particular regions. The pollination time of the trees, grasses, or weeds, the usual amount of each pollen in the air, and the relative ability of these allergens to produce clinical

symptoms should be known. This information can be obtained from reliable biologic supply organizations, botany departments of teaching centers, and allergy clinics.

Skin tests must be applied by the scratch or puncture method. All types of pollen in any given section are used in order to get a complete record of sensitivities. The reactions often reveal just how much the patient will be helped by specific therapy.

If the sensitivity is limited to ragweed and the individual is ill at the time of year when this allergen is most prevalent, pollen treatment should be satisfactory. If the tests show multiple sensitivities to trees, grasses, or weeds, and the patient has symptoms only at the time of weed pollination, the response may still be good.

Should there be minor symptoms throughout the summer with a flareup during the ragweed pollen season, specific therapy will probably not be very satisfactory no matter how carefully the procedure is carried out. For this reason, each hay fever sufferer should have a complete understanding of the value of the treatment. He should not feel that hyposensitization therapy will give perfect results.

Before the specific form of treatment for allergic diseases becomes a highly successful procedure, more must be known concerning the physical and chemical nature of the pollen extracts. In some individuals they act like a true antigen; in others, they fail to make much impression.

This indicates that everything possible must be done to promote a good antibody response. The material injected should be a fresh mixture of the pollen extracts, matching the pa-

tient's sensitivities and the degree of pollination in the community. The inoculations must follow a given schedule closely.

The preseasonal series of subcutaneous injections still tops the list in producing the most satisfactory results. Oral therapy works fairly well in some cases of grass pollen allergy if the preparation is taken in syrup before breakfast.

The coseasonal method in which very small doses of the pollen allergen are given intracutaneously appears to be tolerated by only a few individuals. The practitioner who has not had much instruction and experience concerning this procedure should not attempt it.

Even with the present inadequate

knowledge as to the nature of pollen, can specific therapy be improved? The answer is in the affirmative. An eight-year study of hay fever therapy at the Minneapolis General and University of Minnesota Hospitals revealed that approximately 60% of the failures were due to *lack of cooperation on the part of the patient, the doctor's office, or both in following closely the dosage schedule, to absence of information concerning other allergies of the hay fever sufferer, and to full disregard of special rules of conduct during the hay fever season.*

It is amazing to find how large a number of patients fail to cooperate. If the patient does not keep the appointments, the doctor often makes little or no attempt to have him come



Starting in Practice Doctor?

then get started right
with the **DAILY LOG**

A complete financial record book for physicians. Used by thousands for over 20 years.

As a special introductory offer to doctors just beginning practice, the Dr. Colwell DAILY LOG for 1949 is offered at a reduced rate. You can try this proven system for the remainder of the year—then we believe that, like 90% of our users, you'll reorder for the complete year in 1950. Write for details and descriptive booklet.

COLWELL PUBLISHING COMPANY
239 University Ave. Champaign, Illinois

SPECIAL ARTICLE

regularly. Interruptions are permitted because of reactions to inoculations, although many of these reactions can be overcome by an antihistamine.

Patients with a few clinical symptoms due to tree pollen may be greatly upset by specific ragweed pollen therapy. Too many times the physician then stops the inoculations. It would be better to explain the situation to the patient and try to control side reactions with drugs while continuing the specific treatment.

Some of the poor results are due to the fact that all the sensitivities of the individual are not known. A very young child may be sensitive to foods and an older one may have inhalant allergies.

• The adult may have trouble with

fungi or molds, especially in the southern parts of the United States. Measures to reduce or eliminate the offending foods from the diet or the irritating inhalants from the environment are essential, but this cannot be accomplished in an intelligent way unless a complete set of cutaneous tests has been applied. Lucky is the patient who knows his allergic picture. He can conduct himself accordingly. Such a hay fever sufferer will be more willing to eat properly, rest plenty, play or work less, and avoid exposure to secondary irritants.

COMBINED THERAPY

When specific pollen therapy does not yield full satisfaction in spite of a careful and thorough check of all re-

DOCTOR...

*Lavoris coagulates,
detaches and removes
germ-laden debris,
leaving tissues
cleansed, refreshed
and invigorated.*

ACTIVE INGREDIENTS
Zinc Chloride - Menthol
Formaldehyde - Saccharine
Oil Cinnamon - Oil Cloves
Alcohol 5%

LAVORIS
TRADE MARK
MOUTHWASH AND GARGLE

Distinctive Cleansing Action

DETACHES AND REMOVES GERM HARBORING FILM FROM MOUTH AND THROAT

LAVORIS

THE LAVORIS COMPANY • MINNEAPOLIS 1, MINN.

lated procedures, antihistaminic drugs added to the treatment are sometimes helpful. In these cases, the specific treatment lowers the possibility of complications such as asthma so that the medication can be given in doses small enough to avoid side reactions.



Most distressing is the onset of asthma. This usually appears quite late in the pollen season but may come early. *The presence of asthma usually is an indication for discontinuance of antihistaminic drugs.* There are a few exceptions in which theophorin or hydryllin may be permitted.

The asthma can be mild to moderate in severity and respond well to oral administration of preparations containing racephedrine, propadrine hydrochloride, orthoxine, or ephedrine sulfate. Too often the condition

becomes progressively worse, causing much suffering. Oral medication may then be of little value.

Rectal suppositories containing theophylline or aminophylline may help. The patient can become quite excited and sedation be necessary. Various preparations are available, among which is the T-Bardrin suppository (Angier), containing pentobarbital sodium, phenobarbital sodium, theophylline, ephedrine hydrochloride, and benzocaine in a cocoa butter base.

These suppositories do a good job in many cases but there are some patients who finally require intravenous therapy such as aminophylline in 5% glucose solution. Action of épinephrine is too transient to be of much value in the face of constant exposure to pollen.



TRY ALCO-SPRAY the New, Improved TAPE REMOVER

Gebauers Alco-Spray Removes
Adhesive Tape PAINLESSLY,

- EFFECTIVE!
- EASY TO USE!
- NON-INFLAMMABLE!
- NON-TOXIC!
- NON-IRRITATING!
- ECONOMICAL!

ALCO-SPRAY is now available through your
local surgical supply dealer.

THE GEBAUER CHEMICAL CO.
9410 St. Catherine Ave., Cleveland 4, Ohio

Current Books & Pamphlets

This catalogue is compiled from all available sources, American and foreign, to insure a complete listing of the month's releases.

Medicine

AN ATLAS OF ELECTROCARDIOGRAPHY by William Dressler and Hugo Roesler. 503 pp., ill. Charles C Thomas, Springfield, Ill. \$14

DIE PATHOLOGISCH-ANATOMISCHEN GRUNDLAGEN DER ALLERGIE by W. Eickhoff. 95 pp., ill. Georg Thieme, Stuttgart. 8.40 M.

KLINISCHE INFektionsLEHRE by F. O. Höring. 2d ed. 245 pp. Julius Springer, Berlin. 18 M.

DEGENERATIVE NIERENERKRANKUNGEN by Hansjürgen Oettel. 328 pp. ill. Georg Thieme, Leipzig. 32 M.

DIAGNOSTISCH-THERAPEUTISCHER LEITFADEN DER INNEREN MEDIZIN FÜR DIE ARSTLICHE SPRECHSTUNDE by Felix O. Höring. 259 pp. Ferdinand Enke, Stuttgart. 10.80 M.

THE NOMENCLATURE OF DISEASE. 7th ed. 385 pp. His Majesty's Stationery Office, London. 8s. 6d.

RATIONAL MEDICINE by J. W. Todd. 378 pp. Simpkin Marshall, London. 25s.

Anesthesia

ANAESTHETICS AND THE PATIENT by G. Ostlere. 166 pp. Sigma Books, London. 7s. 6d.

ESTIVIN

Intraocular means of treating HAY FEVER

Prompt, safe symptomatic relief of the distressing hay fever symptoms—sneezing, nasal discharge, eye itching, lacrimation, etc.—is effectively secured by Estivin.

One drop in each eye upon arising, one, before breakfast, and one after breakfast, will usually keep the sufferer comfortable well into the morning. Estivin does not cause drowsiness, or depression, thus permitting application whenever indicated.



Literature and sample on request

Schieffelin & Co.

Pharmaceutical and Research Laboratories
16 Cooper Square • New York 3, N. Y.



CURRENT BOOKS

Ophthalmology

PRACTICAL ORTHOPTICS IN THE TREATMENT OF SQUINT by Thomas Keith Lyle and Sylvia Jackson. 3d ed. 272 pp., ill. H. K. Lewis & Co., London. 35s.

HISTOLOGY AND HISTOPATHOLOGY OF THE EYE AND ITS ADNEXA by I. G. Sommers. 784 pp., ill. Grune & Stratton, New York City. \$12

EINFÜHRUNG IN DIE PHYSIOLOGISCHE OPTIK by Armin von Tschermak-Seysenegg. 2d ed. 213 pp., ill. Springer, Vienna. 45 Sch.

Tuberculosis

PULMONARY TUBERCULOSIS: PATHOLOGY, DIAGNOSIS, MANAGEMENT AND PREVENTION revised by Walter Pagel *et al.* 2d ed. 738 pp., ill. Oxford University Press, New York City. \$18.50

PRESENT CONCEPTS OF REHABILITATION IN TUBERCULOSIS by Norvin C. Kiefer. 398 pp. National Tuberculosis Association, New York City. \$3.50

TUBERCULOSIS OF THE KNEE JOINT by Johannes Mortens. 550 pp. Einar Munksgaard, Copenhagen. 30 Kr.

Orthopedics


ÜBER DIE PRINZIPIEN DER MECHANISCHEN GLIEDERHEILKUNDE by H. Schubjé. 84 pp., ill. Georg Thieme, Leipzig. 8 M.

L'AXE CORPOREL, MUSCULATURE ET INNERVATION: ÉTUDE ANATOMIQUE, PHYSIOLOGIQUE ET PATHOLOGIQUE by André Thomas and J. de Ajuriaguerra. 538 pp., ill. Masson & Co., Paris. 1,650 fr.

TUMORS OF BONE by Charles F. Geschickter and Murray M. Copeland. 3d ed. 810 pp., ill. J. B. Lippincott Co., Philadelphia. \$17.50

Cancer

CANCER, VOLUME II: RADIATIONS, VIRUS, ENVIRONMENT by J. Maisin. 308 pp. Casterman, Tournai-Paris. 120 fr.



Glycerite

of Hydrogen Peroxide *etc* with Carbamide

Instill one-half dropperful into affected ear four times daily

Supplied in one-ounce bottles with dropper

Samples and Literature on request

International Pharmaceutical Corporation

132 Newbury Street, Boston 16, Massachusetts

Constituents:

Hydrogen Peroxide 1.5%

Urea (Carbamide) 2.5%

8 Hydroxyquinoline 0.1%

Dissolved and stabilized in substantially anhydrous glycerol q.s.ad. 30cc.

CURRENT BOOKS

Gynecology

TECHNIQUES CHIRURGICALES VAGINALES: POSSIBILITÉS ET LIMITES by Eric Weber. 144 pp., ill. J. B. Baillière & Fils, Paris. 500 fr.

Psychiatry

SOME COMMON PSYCHOSOMATIC MANIFESTATIONS by J. Barrie Murray. 114 pp. Oxford University Press, New York City. \$2.50

DIAGRAMS OF THE UNCONSCIOUS: HANDWRITING AND PERSONALITY IN MEASUREMENT, EXPERIMENT AND ANALYSIS by Werner Wolff. 423 pp., ill. Grune & Stratton, New York City. \$8

Pharmacology

PENICILLIN AND OTHER ANTIBIOTICS by G. W. S. Andrews and J. Miller. 160 pp., ill. Todd Publishing Group, London. 7s. 6d.

THE CHEMISTRY OF PENICILLIN edited by Hans T. Clarke *et al.* 1,094 pp., ill. Princeton University Press, Princeton, N.J. \$36

HANDBOOK OF MATERIA MEDICA, TOXICOLOGY AND PHARMACOLOGY by Forrest R. Davison. 4th ed. 730 pp., ill. C. V. Mosby Co., St. Louis. \$8.50

THE PLANT ALKALOIDS by Thomas A. Henry. 4th ed. 828 pp. Blakiston Co., Philadelphia. \$12

THE USE OF PENICILLIN AND STREPTOMYCIN by Chester S. Keefer. 72 pp. University of Kansas Press, Lawrence, Kan. \$2

AUREOMYCIN: A NEW ANTIBIOTIC, WITH PAPERS by B. M. Duggar *et al.* 168 pp., ill. New York Academy of Science, New York City. \$2.50

Legal Medicine

TAYLOR'S PRINCIPLES AND PRACTICE OF MEDICAL JURISPRUDENCE, VOLUME II edited by Sir S. Smith. 10th ed. 841 pp. J. & A. Churchill, London. 50s.

20-minute diagnosis of pregnancy



WITH THE
Westerfield®

CHEMICAL
PREGNANCY TEST*
(CARSON-SAEKS METHOD)

94.4% accurate in 431 cases reported¹

FEATURES:

- New simple office procedure—only apparatus required is a beaker and hot plate.
- One or several tests can be run in 20 to 30 minutes.
- No delay—results can be read immediately.

- Economical—costs less per test than other reliable methods.

AVAILABLE: Complete test kit; also refill units and individual replacement parts through your physicians' supply dealer.

1. Ricketts, W. A.; Carson, R. M., and Seeks, R. R. *Am. J. Obst. & Gynec.* 56: 955 (Nov.) 1948.

*U. S. Pat. Applied For.

Westerfield® PHARMACAL CO., INC., DAYTON, OHIO—FINE PHARMACEUTICALS SINCE 1894

CURRENT BOOKS

Venereal Diseases

GLANDES ENDOCRINES ET SYPHILIS: SCHÉMAS CLINIQUES ET THÉRAPEUTIQUES by Paul Blum. 109 pp. Masson & Co., Paris. 250 fr.

Nursing

PRINCIPLES OF PSYCHIATRIC NURSING by Madelene Ingram. 3d ed. 525 pp., ill. W. B. Saunders Co., Philadelphia. \$3.75
EYE, EAR, NOSE AND THROAT MANUAL FOR NURSES by Roy H. Parkinson. 6th ed. 259 pp., ill. C. V. Mosby Co., St. Louis. \$3

Miscellaneous


PHYSICAL ASPECTS OF COLOUR by P. J. Bouma. 312 pp., ill. Elsevier Book Co., New York City. \$5.50
WHAT IS THE LEADING CAUSE OF DEATH? TWO NEW MEASURES by Frank G. Dickinson and Everett L. Walker. 25 pp. American Medical Association, Chicago. Apply

MEDICAL LATIN by Carolyn Lewis. 135 pp. Marshall Jones Co., Francetown, N.H. \$2

GERMAN-ENGLISH DICTIONARY by F. S. Schoenewald. 241 pp. H. K. Lewis & Co., London. 27s. 6d.

DOCTORS OF INFAMY: THE STORY OF THE NAZI MEDICAL CRIMES by Alexander Mitscherlich and Fred Mielke. 172 pp., ill. Henry Schuman, New York City. \$3

DELIVERY ROOM



For 15 Years

GADOMENT has demonstrated a distinctive ability to promote epithelial growth and to stimulate granulation of tissue.

GADOMENT

contains non-destearinated cod liver oil 70%; carbolic acid 0.375%; zinc oxide; benzoin, wax base. Supplied in 1 1/4 oz. and 5 oz. tubes, and 1 lb. jars.

THE E. L. PATCH COMPANY
BOSTON, MASS.

Hospital Swabs

Conform to Federal Specifications GG A-616



Manufactured by Q-TIPS, INC.
Makers of Q-Tips 3" double-tipped swabs
Distributed by
THE SEAMLESS RUBBER COMPANY

ARTHRITIS



ONE CAPSULE WEAPON FOR 2-WAY THERAPY

Many cases of arthritis are accompanied by fibrillitis. Steinberg* showed 143 out of 145 cases of primary fibrillitis aided by high potency vitamin E. EDREX offers both vitamin D and vitamin E. (*An. Int. Med., 19:136-139) Send for free literature.

EDREX VITAMINE PLUS VITAMIN D

WILCO LABORATORIES

800 N. Clark St., Chicago 10, Ill.



**Safe
Removal
of
Skin
Blemishes**

Cauter-izing unit requiring no anesthetic; fully controllable; shockproof. In-terchangeable applicators. A bland technique usable around eyes and hairline.

Cosmo Caution Co.
St. Louis, Mo.

Cosmo Caution Unit

See your dealer or write us

Mechanical Aid in Relief Of Allergic Suffering

As an adjunct to your treatment of inhalant allergic symptoms, recommend the use of the Weaver Nasal Filter. Filters out many possible allergens and irritants from the air breathed. Individually fitted to patient's nostrils by trained technicians. Sanitary silver cups with replaceable filter mats are comfortably worn and are scarcely visible. For literature and address of nearest fitting agency, write

NASAL FILTER CO., Columbus 13, Ohio
Dept. B1

Have You Moved?

If you have changed your address recently, notify us promptly so you will not miss any copies of **MODERN MEDICINE**. Be sure to indicate your old as well as your new address. Send notices to: Circulation Department, **MODERN MEDICINE**, 84 South Tenth Street, Minneapolis 3, Minnesota.

Medical Films

Pertinent information concerning new releases on medical subjects and other recent films that are still available.

New Releases

MEDICINE. The following films available for sale from Castle Films, Division of United World Films Inc., 1445 Park Ave., New York 29:

CLINICAL MALARIA, 26 min.; **EVALUATION OF PHYSICAL FITNESS BY THE STEP-UP TEST**, 12 min.; **SPECIFIC GRAVITY OF THE HEALTHY MAN**, 16 min.; **LIFE CYCLE OF ENDAMOEBIA HISTOLYTICA**, 19 min. silent, color; **MOLES AND MELANOMA**, 7 min. color; **ACTION OF THE EXTRA-OCULAR MUSCLES**, 8 min. color; **PREPARATION AND USE OF HUMAN PLASMA**, 34 min. color; **HEMOLYTIC STREPTOCOCCUS CONTROL**, 13 min.

SURGERY. The following films available for sale from Castle Films, Division of United World Films Inc., 1445 Park Ave., New York 29:

FRACTURE OF THE SURGICAL SPINE: CRUTCHFIELD TONGS REDUCTION 11 min. silent, color; **SKELETAL FIXATION BY THE STADER SPLINT: FRACTURES OF THE TIBIA**, 22 min.; **SKELETAL FIXATION BY THE STADER SPLINT: FRACTURES OF THE OS CALSIS**, 12 min.; **EYE SURGERY: TREATMENT FOR PARESIS OF THE SUPERIOR OBLIQUE**, 7 min. color; **EYE SURGERY: REMOVAL OF INTRAOCULAR FOREIGN BODIES**, 22 min. color; **EYE SURGERY: FIELD MANAGEMENT OF EYE INJURIES**, 20 min. color; **EARLY CARE OF PLASTIC SURGICAL CASES: WOUNDS OF THE HAND**, 14 min. color; **EARLY CARE OF PLASTIC SURGICAL CASES: WOUNDS OF THE FACE AND JAW**, 19 min. color; **AMPUTATIONS: GUILLOTINE AMPUTATIONS OF THE LOWER EXTREMITY**, 12 min. color; **AMPUTATION: REVISION AND REAMPUTATION OF THE STUMP**, 29 min. color; **AMPUTATION: THE UPPER EXTREMITY**, 15 min. color; **PLASTIC SURGERY OF THE HAND**, 12 min. color; **PLASTIC AND RECONSTRUCTION SURGERY OF THE HAND**, 22 min. color; **PLASTIC REPAIR OF THE CHEEK AND LIP**, 10 min. color; **SCIATIC PAIN AND THE INTERVERTEBRAL DISC**, 33 min. color; **SURGICAL MANAGEMENT OF SPINAL CORD INJURIES**, 8 min. color; **FACIAL INJURIES**, 12 min. color; **CARE OF THE FEET**, 10 min. color.

POTENCY



The comprehensive controls under which we make Bayer Aspirin insure uniform potency. In all, over seventy different tests and inspections are employed in making this best-known of all analgesics. Bayer's reputation and acceptance as *the* analgesic for home use is being jealously guarded. In one of America's finest drug plants where Bayer Aspirin is made, *excellence* is the standard.

PATIENTS

... I Have Met

The editors will pay \$1 for each story published. No contribution will be returned. Send your experiences to the Patients I Have Met Editor, MODERN MEDICINE, 84 South Tenth St., Minneapolis 3, Minn.

Adequate Answer

"What," asked the professor, "is the chief difference between male and female?"

"Vas deferens," answered the bright student.—W.H.

"I've got Bright's disease," said the nurse, "and he's got mine."—D.S.

Caveat Emptor

A young man was approached by a lovely lady who said in a hushed voice, "I'm selling."

The young man took one look and sighed, "I'm buying."

A few days later he had a penile drip. Weeks went by. Then the same woman reappeared and again said, "I'm selling."

"What is it now" was the brusque retort, "cancer?"—W.H.



"They're not twins. They just have the same plastic surgeon!"



*"What do you keep in there?
Golf balls?"*

Solar System

After examining a child, I told the mother that a lot of sun was desirable and advised her to sun the baby, front and back, as much as she could. Several days later the mother called to ask if she couldn't sun the baby just in the back. She **was** plum fagged out, she said, from carrying the child to and fro from the backyard to the frontyard and back.

—C.R.S.

The O.B. patient was nearing term and I asked her how she felt. "Doctor," she replied, "it's like building a boat in the basement. As it progresses you wonder how you are going to get it out."

Nothing Common

In checking over reports on surgical specimens for the day the pathologist picked up a rare one, "cervical ass." He called his typist and asked if it was a typographical error.

"Oh no," she answered, "that is what you said."

"What I said," he replied, "was cervical *os*."

"Well, I thought you were being high hat," retorted the girl.—A.G.

*"We have never had other iron salts so efficacious..."*¹

In the treatment of hypochromic anemia the new molybdenum-iron complex, Mol-Iron, is being found to possess *marked advantages* over equivalent dosage of ferrous sulfate alone, or combined with liver-stomach extract or folic acid.

Independent controlled tests^{1, 2, 3} in relatively resistant iron-deficiency anemia of pregnancy have demonstrated a more rapid hemoglobin regeneration as well as greatly improved gastric tolerance.

Investigators have commented: "the increases in hemoglobin (with Mol-Iron) were . . . dramatic and . . . rapid;"¹ there was "more rapid . . . response than with ferrous sulfate . . . ;"² "unusually effective and well tolerated in . . . hypochromic anemia;"³ "...encouraging results obtained with molybdenized ferrous sulfate in the microcytic hypochromic group indicate a better prognosis in these conditions (pregnancy anemia) in the future."⁴

1. Diekmann, W. J., and Priddle, H. D.: *American J. Obstet. & Gynec.* 57:541-546 (March) 1949.
2. Chesley, R. F., and Annitto, J. E.: *Bull. Margaret Hague Maternity Hospital* 1:68-75 (Sept.) 1948.
3. Healy, J. C.: *Journal-Lancet* 66:218-221 (July) 1946.
4. Talso, P. J.: *J. Ins. Med.* 4:31-34 (Dec.-Jan.-Feb.) 1948-1949.

White's

Mol-iron Tablets
Liquid
MOLYBDENIZED FERROUS SULFATE

—a specially processed, co-precipitated, stable complex of molybdenum oxide 3 mg. (1/20 gr.) and ferrous sulfate 195 mg. (3 gr.). Recommended adult dosage: 2 Tablets, t. i. d. Available in bottles of 100 and 1000 tablets and in a highly palatable Liquid, in bottles of 12 fluid ounces.

WHITE LABORATORIES, INC., Pharmaceutical Manufacturers, Newark 7, N. J.




Borchardt MALT SOUP EXTRACT FOR CONSTIPATED BABIES

Borchardt's Malt Soup Extract is a laxative modifier of milk. One or two teaspoonfuls in a single feeding produce a marked change in the stool. Council Accepted. Send for free sample.



BORCHERDT MALT EXTRACT COMPANY
217 N. Wolcott Ave., Chicago 12, Ill.

FOR THE DIABETIC



CELLU

Gelatin Dessert
Answers patients' desert problems. It's sugar-free!

FREE!

32 page catalog of diet foods sent on request.



6 Delicious Flavors

CELLU **LOW CARBOHYDRATE**
Dietary Foods

CHICAGO DIETETIC SUPPLY HOUSE Inc.
1750 West Van Buren Street Chicago 12, Illinois

Evidence accumulates* ...

IN SUPPORT OF T-BARDRIN

(ANGIER)

In a series exceeding 150 cases of bronchial asthma and nasal allergy recently concluded, findings enthusiastically confirm the effectiveness of T-Bardrin therapy through the rectal route.

Highlights of report —

Only the most difficult cases were selected for treatment. Efficacy in children was dramatic with 93% response to medication. With first suppository, symptoms disappeared in a few minutes and most patients remained symptom-free for from one to three days.

Marked response was shown in 75% of adult cases in whom response to other medication was transitory or lacking.

No untoward reactions were noted among the entire series.

*To be published 1949 (author's name on request)

FORMULA

Each T-Bardrin Suppository contains:	
Phenobarbital sodium	0.05 gm.
Phenobarbital sodium	0.05 gm.
(Warning: may be habit forming)	
Theophylline	0.40 gm.
Ephedrine hydrochloride	0.05 gm.
Benzocaine	0.06 gm.
(In a cocoa butter base)	

SAMPLES AND LITERATURE available when requested on professional letterhead

ANGIER CHEMICAL COMPANY

Manufacturing Chemists since 1886

BOSTON 34

MASSACHUSETTS

Optical Illusion

A patient of mine took her bifocal prescription to a dispensing optician. When she got her glasses she asked if the price included \$2.50 for the doctor. Told that it did not, she pointed to the prescription where it said, "Add 2.50." It took some time to explain the meaning of a presbyopic addition.—B.W.



"Did you carve it yourself?"

Compression

"You have the next hour," the English professor told his class, "to write a short story containing religion, sex, royalty, and mystery."

Within a few minutes the pre-med had his story ready: "My God!" exclaimed the king, "The princess is pregnant. Who done it?"—C.A.C.

"How is the patient who swallowed the half dollar?" asked the doctor. "No change yet, sir," replied the nurse.—M.C.

Once Was Enough

The examination of a woman complaining of amenorrhea of short duration was essentially negative. I suggested she could be pregnant, a suggestion she promptly denied. I explained that even though she had been married for several years, pregnancy was a normal possibility.

"Oh no," she answered. "We have a daughter six years old, and we found out what caused that!"—J.D.B.

**TIME-TESTED
MEDICAMENT**



for **CONSTIPATION
HYPERACIDITY**

For more than 75 years, Phillips' Milk of Magnesia has been generally accepted by the medical profession as a standard therapeutic agent for constipation and gastric hyperacidity.

As a laxative—Phillips' mild, yet thorough action is ideal for both adults and children.

As an antacid—Phillips' affords fast, effective relief. Contains no carbonates, hence produces no discomforting flatulence.

DOSAGE: { Laxative: 2 to 4 tablespoonfuls
Antacid: 1 to 4 teaspoonfuls, or
1 to 4 tablets



PHILLIPS' MILK OF MAGNESIA

Prepared only by **THE CHAS. H. PHILLIPS CO. DIVISION**, 170 Varick Street, New York 13, N. Y.
of Sterling Drug Inc.

Domeboro Powder
MAKES A MODERNIZED BUROW'S SOLUTION
 1:20
 in calculated dose packet

New

Sample on Request

DOME CHEMICALS, INC. 250 E. 43rd STREET
 NEW YORK 17, N.Y.

Your Kind of Answers:
 All Reviewers Acclaim
Dr. Leonard H. Biskind's
"HAVING YOUR BABY"
 MODERN INSTRUCTIONS FOR
 EXPECTANT MOTHERS"

Sound. Understandable. Thorough. Up-to-the-minute. Indispensable in a busy practice.

AND—INEXPENSIVE

\$1 per copy; 10 copies less 10%; 25 copies less 20%; 50 or more copies less 25%. Order today.

Western Journal of Surgery
 Publishing Company
 202 PHOENIX BUILDING, PORTLAND 4, OREGON

UROPATHY

Relief
Sedation
Bacteriostasis

SEDURIN

Formula / Fluid oz.
 Methenamine . . 18 gr.
 Sandalwood . . 30 gr.
 Saw Palmetto . . 30 gr.
 Zea 30 gr.

Alcohol 9%

Available on prescription only, in 8-oz. bottles.

FREE SAMPLE

DRUG SPECIALTIES, INC.
 218 Boyd Street, Los Angeles 54, Calif.

Professional Sample, Please:

M. D.

INDEX TO ADVERTISERS

The publishers are not responsible for any errors or omissions.

Abbott Laboratories	5
Amherst Research, Inc.	22
Angier Chemical Co.	104
Armour Laboratories, The	82-83
Barnes, A. C., Company	13
Bausch & Lomb Optical Co.	32
Bayer Aspirin	101
Belmont Laboratories	27
Birtcher Corporation, The	28
Borchardt Malt Extract Company	104
Borden Company, The	56-57
Burroughs Wellcome & Co. (U.S.A.)	29
Chicago Dietetic Supply House, Inc.	104
Ciba Pharmaceutical Products, Inc.	4th Cover
Colwell Publishing Co.	93
Commercial Solvents Corporation	80
Cosmo Cautery Co.	106
Dome Chemicals, Inc.	100
Drew Pharmacal Co., Inc.	106
Drug Specialties, Inc.	106
Eastman Kodak Company	76-77
Eaton Laboratories, Inc.	75
Endo	108
Gebauer Chemical Company, The	95
Hoffmann-La Roche, Inc.	11
International Pharmaceutical Corporation	97
Johnson & Johnson	87
Lavoris	94
Lederle Laboratories, Inc.	45
McNeil Laboratories	3
Merrell, The Wm. S. Company	2nd Cover
Nasal Filter Company	100
Parke, Davis & Company	41
Patch, The E. L., Company	99
Phillips Co., The Chas. H.	105
Pitman Moore Company	15
Professional Printing Company, Inc.	7
Ritter Co., Inc.	79
Robins, A. H., Company, Inc.	16-17, 35, 36
Roebig, J. B., & Company	8-9
Schering Corporation	23
Schiffelin & Co.	96
Seamless Rubber Co., The	99
Sharp & Dohme	33
Sherman Laboratories	19
Shield Laboratories	4
Smith, Kline & French Laboratories	21
Smith, Martin H., Company	89
Special Milk Products, Inc.	26
Strauss Laboratories	39
Ulmer Pharmacal Company	107
Valsey-Bristol Shoe Co., Inc.	20
Walker, Myron L., Co., Inc.	30
Walker Vitamin Products, Inc.	38
Warren-Teed Products Company, The	25
Westernfield Pharmacal Co., Inc.	98
Western Journal of Surgery Publishing Co.	106
White Laboratories, Inc.	31, 103
Whitehall Pharmacal Company	37
Whittier Laboratories	42-43, 91
Wilco Laboratories	100
Winthrop-Stearns, Inc.	3rd Cover
Wyeth Incorporated	46

Immediate and Continued Dominance Over Syphilis

Is the purpose of every logical plan for the treatment of this infection

To fulfill this purpose, the chosen agent must be a spirocheticide of proved power, without danger in its proper application and of such chemical structure that release of its essential element into the tissues begins at once and continues well into the period of the following injections. It must enable the organism to build up with it a considerable reserve of this essential element, to maintain an unbroken spirocheticidal effect, to create the necessary serological action, and to cause the Wassermann to remain permanently negative.

In the Liposoluble bismuth preparation: "Biliposol" are to be found these desirable properties

BILIPOSOL

TRADE
MARK

REG. U. S.
PAT. OFF.

*A Special Lipoid Soluble Bismuth in Clear
Homogeneous Solution for the Treatment of*

SYPHILIS

**INDICATED FOR THE SUCCESSFUL TREATMENT
OF ALL STAGES OF SYPHILIS**

Its prompt and complete absorption insures immediate action and total effectiveness.

**BILIPOSOL REQUIRES NO HOSPITALIZATION AND IS A MOST SATISFACTORY
TREATMENT OF PRIVATE OFFICE PATIENTS, AS THERE IS NO PAIN FROM PROPER
INJECTION, NO TOXICITY OR SHOCK.**

"It can be said that, as a whole, the cases of primary or secondary syphilis treated during the first few months of the infection have a negative Wassermann in at least 75% of cases 2 months after the beginning of the treatment, and this immediate action seems to be equally a lasting one in the majority of cases."

—Dr. C. Levaditi.

BOXES OF 12, 50 AND 100 — 2 C.C. AMPOULES — EACH CONTAINING 8 CTG. BISMUTH

ORDER TODAY

IF YOUR DRUGGIST CANNOT SUPPLY YOU—WE WILL

ULMER LABORATORIES

414 So. Sixth Street

Minneapolis, Minnesota

Sole General Distributors for United States and Canada

Controlled action in digestive distress

When pain, heartburn, belching, nausea, or unstable colon are due to gastrointestinal spasm, Mesopin provides an effective means for prompt relief. Its selective antispasmodic action on the digestive tract controls spasticity with virtual freedom from the undesirable side effects of atropine or belladonna. Thus, symptomatic relief of many spastic disturbances of the stomach or intestines can be achieved with discrimination and greater safety.

Mesopin*


selective
gastrointestinal
antispasmodic



Supplied: Mesopin (2.5 mg. per tablet)
is available on prescription in
bottles of 100 tablets.
Samples sent on request.

*brand of homatropine methyl bromide

Endo Products Inc., Richmond Hill 10, N.Y.



Specific
for
Trichomonas

Trichomonas vaginalis, the most common cause of leukorrhea, is readily eradicated by Devegan. Furthermore, Devegan restores the normal vaginal flora and pH level, relieves irritation, and eliminates the offensive leukorrheal discharge. Powder for insufflation at the office (10 Gm. vials, 1 oz. and 8 oz. bottles). Tablets for patient's use at home (boxes of 25 and 250).

EFFECTIVE COMBINATION TREATMENT FOR LEUKORRHEA



Winthrop Stearns INC.
NEW YORK 13, N. Y. WINDSOR, ONT.

DEVEGAN, trademark registered

NEW METHOD FOR RELIEF OF ALLERGIC NASAL CONGESTION

Distributes mist of minute droplets of Pyr-
benzamine hydrochloride Nasal Solution 0.5%
throughout nasal passages.

Relief is immediate—complete—prolonged.
No side reactions except occasional transient
stinging. Convenient to use and carry.

Non-refillable. Provides several hundred
applications. Dosage one application to each
nostril every 3 to 4 hours.



PHARMACEUTICAL PRODUCTS, INC.
SUMMIT, NEW JERSEY

PYR-BENZAMINE (brand of triphenylamine)
U.S. Pat. 2,711,111